

表 3137 nPCR(g/kg/day) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	8	300	2,977	6,289	3,149	561	63	6	1	5	13,359	767	14,126	0.91	0.17
(%)	(0.1)	(2.2)	(22.3)	(47.1)	(23.6)	(4.2)	(0.5)	(0.0)	(0.0)	(0.0)	(100.0)				
慢性腎盂腎炎	1	11	100	169	79	15	4				379	19	398	0.90	0.18
(%)	(0.3)	(2.9)	(26.4)	(44.6)	(20.8)	(4.0)	(1.1)				(100.0)				
急速進行性糸球体腎炎	1	7	41	110	62	14	3	1			239	11	250	0.92	0.19
(%)	(0.4)	(2.9)	(17.2)	(46.0)	(25.9)	(5.9)	(1.3)	(0.4)			(100.0)				
妊娠腎/妊娠中毒症	1	8	64	147	97	15	2				334	14	348	0.93	0.17
(%)	(0.3)	(2.4)	(19.2)	(44.0)	(29.0)	(4.5)	(0.6)				(100.0)				
その他の分類不能の腎炎		4	47	77	39	11					178	8	186	0.90	0.17
(%)		(2.2)	(26.4)	(43.3)	(21.9)	(6.2)					(100.0)				
多発性嚢胞腎		29	328	625	223	34	2				1,241	89	1,330	0.88	0.15
(%)		(2.3)	(26.4)	(50.4)	(18.0)	(2.7)	(0.2)				(100.0)				
腎硬化症	4	90	647	943	387	66	11				2,148	104	2,252	0.87	0.17
(%)	(0.2)	(4.2)	(30.1)	(43.9)	(18.0)	(3.1)	(0.5)				(100.0)				
悪性高血圧	1	5	67	122	48	15	1				259	11	270	0.89	0.17
(%)	(0.4)	(1.9)	(25.9)	(47.1)	(18.5)	(5.8)	(0.4)				(100.0)				
糖尿病性腎症	19	542	3,644	4,291	1,451	211	16	5			10,179	666	10,845	0.84	0.16
(%)	(0.2)	(5.3)	(35.8)	(42.2)	(14.3)	(2.1)	(0.2)	(0.0)			(100.0)				
SLE腎炎		4	68	131	84	25	5				317	20	337	0.94	0.19
(%)		(1.3)	(21.5)	(41.3)	(26.5)	(7.9)	(1.6)				(100.0)				
アミロイド腎		1	15	15	8	3					42	1	43	0.89	0.18
(%)		(2.4)	(35.7)	(35.7)	(19.0)	(7.1)					(100.0)				
痛風腎	1	2	37	61	23	7					131	11	142	0.87	0.17
(%)	(0.8)	(1.5)	(28.2)	(46.6)	(17.6)	(5.3)					(100.0)				
先天性代謝異常による腎不全			7	16	9	3					35	6	41	0.95	0.16
(%)			(20.0)	(45.7)	(25.7)	(8.6)					(100.0)				
腎・尿路結核		1	6	6	3	3					19	1	20	0.90	0.22
(%)		(5.3)	(31.6)	(31.6)	(15.8)	(15.8)					(100.0)				
腎・尿路結石		2	24	25	9	3					63	1	64	0.86	0.17
(%)		(3.2)	(38.1)	(39.7)	(14.3)	(4.8)					(100.0)				
腎・尿路腫瘍		1	15	34	8	1					59	3	62	0.88	0.13
(%)		(1.7)	(25.4)	(57.6)	(13.6)	(1.7)					(100.0)				
閉塞性尿路障害		1	13	31	16	3					64	5	69	0.91	0.16
(%)		(1.6)	(20.3)	(48.4)	(25.0)	(4.7)					(100.0)				
骨髄腫			6	3	2						11		11	0.82	0.13
(%)			(54.5)	(27.3)	(18.2)						(100.0)				
腎形成不全			11	46	24	8	2				91	4	95	0.97	0.18
(%)			(12.1)	(50.5)	(26.4)	(8.8)	(2.2)				(100.0)				
不明	5	92	732	1,231	551	115	12			1	2,739	215	2,954	0.89	0.18
(%)	(0.2)	(3.4)	(26.7)	(44.9)	(20.1)	(4.2)	(0.4)			(0.0)	(100.0)				
再導入	1	6	64	172	100	23	2				368	17	385	0.94	0.17
(%)	(0.3)	(1.6)	(17.4)	(46.7)	(27.2)	(6.3)	(0.5)				(100.0)				
その他	1	24	160	336	159	31	5	1			717	26	743	0.90	0.17
(%)	(0.1)	(3.3)	(22.3)	(46.9)	(22.2)	(4.3)	(0.7)	(0.1)			(100.0)				
合計	43	1,130	9,073	14,880	6,531	1,167	128	13	1	6	32,972	1,999	34,971	0.88	0.17
(%)	(0.1)	(3.4)	(27.5)	(45.1)	(19.8)	(3.5)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし															
(%)															
総計	43	1,130	9,073	14,880	6,531	1,167	128	13	1	6	32,972	1,999	34,971	0.88	0.17
(%)	(0.1)	(3.4)	(27.5)	(45.1)	(19.8)	(3.5)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査