

表 3144 nPCR(g/kg/day) %クレアチニン産生速度(%)別 (血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
60未満	12	83	228	216	78	10	5				632		632	0.80	0.20
(%)	(1.9)	(13.1)	(36.1)	(34.2)	(12.3)	(1.6)	(0.8)				(100.0)				
60～	1	57	200	183	61	14	2				518		518	0.81	0.18
(%)	(0.2)	(11.0)	(38.6)	(35.3)	(11.8)	(2.7)	(0.4)				(100.0)				
70～	1	51	333	349	142	27	3				907		907	0.84	0.18
(%)	(0.1)	(5.6)	(36.7)	(38.5)	(15.7)	(3.0)	(0.3)				(100.0)				
80～	5	61	429	555	230	50	8	1			1,339		1,339	0.86	0.18
(%)	(0.4)	(4.6)	(32.0)	(41.4)	(17.2)	(3.7)	(0.6)	(0.1)			(100.0)				
90～	1	61	498	823	414	87	6	2			1,892		1,892	0.89	0.17
(%)	(0.1)	(3.2)	(26.3)	(43.5)	(21.9)	(4.6)	(0.3)	(0.1)			(100.0)				
100～	2	43	507	1,017	544	112	14	3			2,242		2,242	0.91	0.17
(%)	(0.1)	(1.9)	(22.6)	(45.4)	(24.3)	(5.0)	(0.6)	(0.1)			(100.0)				
110～		23	379	929	580	116	19	1			2,047		2,047	0.94	0.17
(%)		(1.1)	(18.5)	(45.4)	(28.3)	(5.7)	(0.9)	(0.0)			(100.0)				
120～	1	14	234	724	439	103	6	1			1,522		1,522	0.95	0.16
(%)	(0.1)	(0.9)	(15.4)	(47.6)	(28.8)	(6.8)	(0.4)	(0.1)			(100.0)				
130～		3	116	373	276	51	10				829		829	0.96	0.16
(%)		(0.4)	(14.0)	(45.0)	(33.3)	(6.2)	(1.2)				(100.0)				
140～		6	70	205	154	28	5				468		468	0.96	0.16
(%)		(1.3)	(15.0)	(43.8)	(32.9)	(6.0)	(1.1)				(100.0)				
合計	23	402	2,994	5,374	2,918	598	78	8		1	12,396		12,396	0.90	0.18
(%)	(0.2)	(3.2)	(24.2)	(43.4)	(23.5)	(4.8)	(0.6)	(0.1)		(0.0)	(100.0)				
記載なし				1			2		1	2	6	793	799	1.88	0.75
(%)				(16.7)			(33.3)		(16.7)	(33.3)	(100.0)				
総計	23	402	2,994	5,375	2,918	598	80	8	1	3	12,402	793	13,195	0.90	0.18
(%)	(0.2)	(3.2)	(24.1)	(43.3)	(23.5)	(4.8)	(0.6)	(0.1)	(0.0)	(0.0)	(100.0)				
平均	65.22	80.79	95.14	103.99	108.43	108.62	106.83	104.11		79.80	102.31		102.31		
標準偏差	25.62	25.46	23.90	22.55	22.72	21.26	28.20	12.49			23.89		23.89		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査