

表 3146 nPCR(g/kg/day) 透析前CRP濃度(mg/dL)別(血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前CRP濃度	0.4未満	0.4～	0.6～	0.8～	1.0～	1.2～	1.4～	1.6～	1.8～	2.0～	合計	記載なし	総計	平均	標準偏差
0.2未満	16	425	4,412	8,163	3,741	698	68	6		4	17,533	548	18,081	0.90	0.17
(%)	(0.1)	(2.4)	(25.2)	(46.6)	(21.3)	(4.0)	(0.4)	(0.0)		(0.0)	(100.0)				
0.2～	6	222	1,649	2,443	1,003	173	23	3		1	5,523	207	5,730	0.87	0.17
(%)	(0.1)	(4.0)	(29.9)	(44.2)	(18.2)	(3.1)	(0.4)	(0.1)		(0.0)	(100.0)				
0.5～	3	145	814	1,167	467	77	8	1	1		2,683	87	2,770	0.86	0.17
(%)	(0.1)	(5.4)	(30.3)	(43.5)	(17.4)	(2.9)	(0.3)	(0.0)	(0.0)		(100.0)				
1.0～	3	101	595	705	260	39	6	2			1,711	55	1,766	0.84	0.17
(%)	(0.2)	(5.9)	(34.8)	(41.2)	(15.2)	(2.3)	(0.4)	(0.1)			(100.0)				
2.0～	5	81	334	371	145	21	5				962	45	1,007	0.83	0.18
(%)	(0.5)	(8.4)	(34.7)	(38.6)	(15.1)	(2.2)	(0.5)				(100.0)				
4.0～	1	24	121	128	54	9	1				338	17	355	0.84	0.18
(%)	(0.3)	(7.1)	(35.8)	(37.9)	(16.0)	(2.7)	(0.3)				(100.0)				
6.0～	1	7	48	54	25	7	1				143	7	150	0.85	0.20
(%)	(0.7)	(4.9)	(33.6)	(37.8)	(17.5)	(4.9)	(0.7)				(100.0)				
8.0～		4	23	37	11						75	10	85	0.84	0.15
(%)		(5.3)	(30.7)	(49.3)	(14.7)						(100.0)				
10.0～		9	35	25	11	1					81	10	91	0.80	0.17
(%)		(11.1)	(43.2)	(30.9)	(13.6)	(1.2)					(100.0)				
15.0～	1	6	27	20	14	2					70	5	75	0.83	0.19
(%)	(1.4)	(8.6)	(38.6)	(28.6)	(20.0)	(2.9)					(100.0)				
合計	36	1,024	8,058	13,113	5,731	1,027	112	12	1	5	29,119	991	30,110	0.88	0.17
(%)	(0.1)	(3.5)	(27.7)	(45.0)	(19.7)	(3.5)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
記載なし	7	106	1,015	1,767	800	140	16	1		1	3,853	1,008	4,861	0.89	0.17
(%)	(0.2)	(2.8)	(26.3)	(45.9)	(20.8)	(3.6)	(0.4)	(0.0)		(0.0)	(100.0)				
総計	43	1,130	9,073	14,880	6,531	1,167	128	13	1	6	32,972	1,999	34,971	0.88	0.17
(%)	(0.1)	(3.4)	(27.5)	(45.1)	(19.8)	(3.5)	(0.4)	(0.0)	(0.0)	(0.0)	(100.0)				
平均	1.55	1.01	0.66	0.48	0.45	0.46	0.46	0.46	0.69	0.11	0.54	0.83	0.55		
標準偏差	3.41	2.52	1.87	1.51	1.46	2.75	1.02	0.59		0.16	1.71	2.40	1.74		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査