

表 3159 %クレアチニン産生速度(%) 年齢(歳)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

年齢	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
15歳未満	3	1	1								5	1	6	54.94	21.50
(%)	(60.0)	(20.0)	(20.0)								(100.0)				
15歳～	28	21	32	41	62	94	80	27	13	7	405	36	441	98.05	23.91
(%)	(6.9)	(5.2)	(7.9)	(10.1)	(15.3)	(23.2)	(19.8)	(6.7)	(3.2)	(1.7)	(100.0)				
30歳～	312	247	460	802	1,129	1,372	1,226	839	373	154	6,914	566	7,480	101.67	22.33
(%)	(4.5)	(3.6)	(6.7)	(11.6)	(16.3)	(19.8)	(17.7)	(12.1)	(5.4)	(2.2)	(100.0)				
45歳～	1,513	1,339	2,271	3,537	4,622	5,069	4,775	3,354	1,720	867	29,067	2,371	31,438	100.68	23.65
(%)	(5.2)	(4.6)	(7.8)	(12.2)	(15.9)	(17.4)	(16.4)	(11.5)	(5.9)	(3.0)	(100.0)				
60歳～	5,249	4,068	6,456	8,960	11,536	12,607	11,381	8,514	4,852	3,313	76,936	6,647	83,583	99.91	25.58
(%)	(6.8)	(5.3)	(8.4)	(11.6)	(15.0)	(16.4)	(14.8)	(11.1)	(6.3)	(4.3)	(100.0)				
75歳～	5,028	3,207	4,433	5,998	7,132	7,096	6,100	4,396	2,537	1,993	47,920	4,292	52,212	95.72	27.74
(%)	(10.5)	(6.7)	(9.3)	(12.5)	(14.9)	(14.8)	(12.7)	(9.2)	(5.3)	(4.2)	(100.0)				
90歳～	417	242	301	345	377	357	292	208	113	115	2,767	278	3,045	90.90	30.14
(%)	(15.1)	(8.7)	(10.9)	(12.5)	(13.6)	(12.9)	(10.6)	(7.5)	(4.1)	(4.2)	(100.0)				
合計	12,550	9,125	13,954	19,683	24,858	26,595	23,854	17,338	9,608	6,449	164,014	14,191	178,205	98.74	25.96
(%)	(7.7)	(5.6)	(8.5)	(12.0)	(15.2)	(16.2)	(14.5)	(10.6)	(5.9)	(3.9)	(100.0)				
不明												4	4		
(%)															
記載なし															
(%)															
総計	12,550	9,125	13,954	19,683	24,858	26,595	23,854	17,338	9,608	6,449	164,014	14,195	178,209	98.74	25.96
(%)	(7.7)	(5.6)	(8.5)	(12.0)	(15.2)	(16.2)	(14.5)	(10.6)	(5.9)	(3.9)	(100.0)				
平均	71.30	69.95	68.80	68.02	67.44	66.76	66.47	66.70	67.42	69.43	67.81	68.28	67.85		
標準偏差	11.90	11.96	12.01	12.21	12.25	12.36	12.27	11.92	11.48	10.89	12.15	12.14	12.15		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査