

表 3161 %クレアチニン産生速度(%) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

原疾患	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,541	1,915	3,204	5,003	7,441	9,167	9,630	7,798	4,691	3,202	54,592	4,855	59,447	105.66	24.62
(%)	(4.7)	(3.5)	(5.9)	(9.2)	(13.6)	(16.8)	(17.6)	(14.3)	(8.6)	(5.9)	(100.0)				
慢性腎盂腎炎	105	85	127	196	250	302	258	219	97	65	1,704	113	1,817	100.72	25.29
(%)	(6.2)	(5.0)	(7.5)	(11.5)	(14.7)	(17.7)	(15.1)	(12.9)	(5.7)	(3.8)	(100.0)				
急速進行性糸球体腎炎	151	90	104	112	150	173	163	122	78	49	1,192	89	1,281	95.56	30.31
(%)	(12.7)	(7.6)	(8.7)	(9.4)	(12.6)	(14.5)	(13.7)	(10.2)	(6.5)	(4.1)	(100.0)				
妊娠腎/妊娠中毒症	30	26	61	80	113	163	183	135	80	53	924	75	999	106.76	23.04
(%)	(3.2)	(2.8)	(6.6)	(8.7)	(12.2)	(17.6)	(19.8)	(14.6)	(8.7)	(5.7)	(100.0)				
その他の分類不能の腎炎	46	27	40	64	97	125	132	89	47	37	704	59	763	102.93	26.33
(%)	(6.5)	(3.8)	(5.7)	(9.1)	(13.8)	(17.8)	(18.8)	(12.6)	(6.7)	(5.3)	(100.0)				
多発性嚢胞腎	224	244	428	773	1,040	1,275	1,087	685	337	186	6,279	505	6,784	102.01	21.81
(%)	(3.6)	(3.9)	(6.8)	(12.3)	(16.6)	(20.3)	(17.3)	(10.9)	(5.4)	(3.0)	(100.0)				
腎硬化症	997	716	1,104	1,484	1,984	2,149	1,976	1,561	945	707	13,623	1,054	14,677	100.44	26.70
(%)	(7.3)	(5.3)	(8.1)	(10.9)	(14.6)	(15.8)	(14.5)	(11.5)	(6.9)	(5.2)	(100.0)				
悪性高血圧	107	50	103	135	165	217	234	181	99	65	1,356	109	1,465	101.84	27.15
(%)	(7.9)	(3.7)	(7.6)	(10.0)	(12.2)	(16.0)	(17.3)	(13.3)	(7.3)	(4.8)	(100.0)				
糖尿病性腎症	6,375	4,730	7,097	9,391	10,519	9,472	6,918	4,041	1,923	1,094	61,560	5,290	66,850	91.64	24.87
(%)	(10.4)	(7.7)	(11.5)	(15.3)	(17.1)	(15.4)	(11.2)	(6.6)	(3.1)	(1.8)	(100.0)				
SLE腎炎	104	76	112	155	170	189	146	108	45	22	1,127	88	1,215	94.72	25.22
(%)	(9.2)	(6.7)	(9.9)	(13.8)	(15.1)	(16.8)	(13.0)	(9.6)	(4.0)	(2.0)	(100.0)				
アミロイド腎	39	17	21	28	28	30	17	8	10	2	200	30	230	86.10	28.04
(%)	(19.5)	(8.5)	(10.5)	(14.0)	(14.0)	(15.0)	(8.5)	(4.0)	(5.0)	(1.0)	(100.0)				
痛風腎	23	27	36	76	92	106	109	89	46	45	649	56	705	105.53	24.19
(%)	(3.5)	(4.2)	(5.5)	(11.7)	(14.2)	(16.3)	(16.8)	(13.7)	(7.1)	(6.9)	(100.0)				
先天性代謝異常による腎不全	13	8	11	16	26	26	19	16	2	4	141	13	154	96.41	24.19
(%)	(9.2)	(5.7)	(7.8)	(11.3)	(18.4)	(18.4)	(13.5)	(11.3)	(1.4)	(2.8)	(100.0)				
腎・尿路結核	14	6	6	13	16	26	24	10	8	2	125	10	135	97.03	26.93
(%)	(11.2)	(4.8)	(4.8)	(10.4)	(12.8)	(20.8)	(19.2)	(8.0)	(6.4)	(1.6)	(100.0)				
腎・尿路結石	13	20	22	36	48	57	44	42	27	24	333	21	354	104.18	24.96
(%)	(3.9)	(6.0)	(6.6)	(10.8)	(14.4)	(17.1)	(13.2)	(12.6)	(8.1)	(7.2)	(100.0)				
腎・尿路腫瘍	36	19	20	41	46	75	73	58	40	33	441	32	473	104.46	28.50
(%)	(8.2)	(4.3)	(4.5)	(9.3)	(10.4)	(17.0)	(16.6)	(13.2)	(9.1)	(7.5)	(100.0)				
閉塞性尿路障害	55	26	29	49	54	58	55	24	21	11	382	32	414	92.18	29.68
(%)	(14.4)	(6.8)	(7.6)	(12.8)	(14.1)	(15.2)	(14.4)	(6.3)	(5.5)	(2.9)	(100.0)				
骨髄腫	11	8	12	12	7	10	6	6	3	2	77	11	88	87.82	28.26
(%)	(14.3)	(10.4)	(15.6)	(15.6)	(9.1)	(13.0)	(7.8)	(7.8)	(3.9)	(2.6)	(100.0)				
腎形成不全	9	14	17	36	58	60	72	45	15	7	333	26	359	103.95	20.92
(%)	(2.7)	(4.2)	(5.1)	(10.8)	(17.4)	(18.0)	(21.6)	(13.5)	(4.5)	(2.1)	(100.0)				
不明	1,210	762	1,019	1,457	1,855	2,146	2,014	1,552	823	669	13,507	1,323	14,830	99.19	27.65
(%)	(9.0)	(5.6)	(7.5)	(10.8)	(13.7)	(15.9)	(14.9)	(11.5)	(6.1)	(5.0)	(100.0)				
再導入	47	50	81	107	157	204	183	157	63	28	1,077	105	1,182	102.36	22.96
(%)	(4.4)	(4.6)	(7.5)	(9.9)	(14.6)	(18.9)	(17.0)	(14.6)	(5.8)	(2.6)	(100.0)				
その他	400	209	300	419	542	565	511	392	208	142	3,688	299	3,987	96.57	28.52
(%)	(10.8)	(5.7)	(8.1)	(11.4)	(14.7)	(15.3)	(13.9)	(10.6)	(5.6)	(3.9)	(100.0)				
合計	12,550	9,125	13,954	19,683	24,858	26,595	23,854	17,338	9,608	6,449	164,014	14,195	178,209	98.74	25.96
(%)	(7.7)	(5.6)	(8.5)	(12.0)	(15.2)	(16.2)	(14.5)	(10.6)	(5.9)	(3.9)	(100.0)				
記載なし															
(%)															
総計	12,550	9,125	13,954	19,683	24,858	26,595	23,854	17,338	9,608	6,449	164,014	14,195	178,209	98.74	25.96
(%)	(7.7)	(5.6)	(8.5)	(12.0)	(15.2)	(16.2)	(14.5)	(10.6)	(5.9)	(3.9)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査