

表 3169 %クレアチニン産生速度(%) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
3未満	1,945	680	744	728	725	604	425	237	107	77	6,272	617	6,889	76.22	30.17
(%)	(31.0)	(10.8)	(11.9)	(11.6)	(11.6)	(9.6)	(6.8)	(3.8)	(1.7)	(1.2)	(100.0)				
3.0～	1,410	696	861	1,031	1,095	1,004	740	468	238	129	7,672	564	8,236	86.17	28.64
(%)	(18.4)	(9.1)	(11.2)	(13.4)	(14.3)	(13.1)	(9.6)	(6.1)	(3.1)	(1.7)	(100.0)				
3.5～	1,835	1,049	1,405	1,723	1,956	1,908	1,554	1,011	572	303	13,316	878	14,194	91.00	27.88
(%)	(13.8)	(7.9)	(10.6)	(12.9)	(14.7)	(14.3)	(11.7)	(7.6)	(4.3)	(2.3)	(100.0)				
4.0～	2,016	1,386	1,978	2,648	3,143	3,124	2,680	1,759	913	572	20,219	1,233	21,452	94.77	26.19
(%)	(10.0)	(6.9)	(9.8)	(13.1)	(15.5)	(15.5)	(13.3)	(8.7)	(4.5)	(2.8)	(100.0)				
4.5～	1,711	1,411	2,223	3,049	3,787	3,933	3,608	2,629	1,386	959	24,696	1,382	26,078	98.91	25.11
(%)	(6.9)	(5.7)	(9.0)	(12.3)	(15.3)	(15.9)	(14.6)	(10.6)	(5.6)	(3.9)	(100.0)				
5.0～	1,321	1,274	2,057	3,092	3,917	4,278	3,807	2,908	1,516	1,066	25,236	1,388	26,624	101.02	24.35
(%)	(5.2)	(5.0)	(8.2)	(12.3)	(15.5)	(17.0)	(15.1)	(11.5)	(6.0)	(4.2)	(100.0)				
5.5～	891	983	1,693	2,532	3,349	3,844	3,495	2,735	1,521	995	22,038	1,084	23,122	102.91	23.56
(%)	(4.0)	(4.5)	(7.7)	(11.5)	(15.2)	(17.4)	(15.9)	(12.4)	(6.9)	(4.5)	(100.0)				
6.0～	110	158	252	437	590	663	566	479	284	208	3,747	219	3,966	104.48	23.22
(%)	(2.9)	(4.2)	(6.7)	(11.7)	(15.7)	(17.7)	(15.1)	(12.8)	(7.6)	(5.6)	(100.0)				
6.1～	439	457	882	1,361	1,963	2,247	2,055	1,548	909	659	12,520	645	13,165	104.29	23.25
(%)	(3.5)	(3.7)	(7.0)	(10.9)	(15.7)	(17.9)	(16.4)	(12.4)	(7.3)	(5.3)	(100.0)				
6.5～	302	382	744	1,122	1,654	1,875	1,938	1,394	810	558	10,779	583	11,362	105.25	22.86
(%)	(2.8)	(3.5)	(6.9)	(10.4)	(15.3)	(17.4)	(18.0)	(12.9)	(7.5)	(5.2)	(100.0)				
7.0～	179	237	406	759	1,015	1,202	1,159	818	542	373	6,690	390	7,080	105.54	22.86
(%)	(2.7)	(3.5)	(6.1)	(11.3)	(15.2)	(18.0)	(17.3)	(12.2)	(8.1)	(5.6)	(100.0)				
7.5～	124	138	241	447	590	703	702	526	313	189	3,973	225	4,198	105.09	22.65
(%)	(3.1)	(3.5)	(6.1)	(11.3)	(14.9)	(17.7)	(17.7)	(13.2)	(7.9)	(4.8)	(100.0)				
8.0～	75	87	154	242	365	437	390	295	176	137	2,358	127	2,485	105.15	23.20
(%)	(3.2)	(3.7)	(6.5)	(10.3)	(15.5)	(18.5)	(16.5)	(12.5)	(7.5)	(5.8)	(100.0)				
8.5～	57	51	110	157	243	265	276	175	116	81	1,531	70	1,601	104.37	23.75
(%)	(3.7)	(3.3)	(7.2)	(10.3)	(15.9)	(17.3)	(18.0)	(11.4)	(7.6)	(5.3)	(100.0)				
9.0～	88	101	162	273	386	428	384	287	167	120	2,396	141	2,537	103.71	23.96
(%)	(3.7)	(4.2)	(6.8)	(11.4)	(16.1)	(17.9)	(16.0)	(12.0)	(7.0)	(5.0)	(100.0)				
合計	12,503	9,090	13,912	19,601	24,778	26,515	23,779	17,269	9,570	6,426	163,443	9,546	172,989	98.74	25.95
(%)	(7.6)	(5.6)	(8.5)	(12.0)	(15.2)	(16.2)	(14.5)	(10.6)	(5.9)	(3.9)	(100.0)				
記載なし	47	35	42	82	80	80	75	69	38	23	571	4,649	5,220	98.65	27.14
(%)	(8.2)	(6.1)	(7.4)	(14.4)	(14.0)	(14.0)	(13.1)	(12.1)	(6.7)	(4.0)	(100.0)				
総計	12,550	9,125	13,954	19,683	24,858	26,595	23,854	17,338	9,608	6,449	164,014	14,195	178,209	98.74	25.96
(%)	(7.7)	(5.6)	(8.5)	(12.0)	(15.2)	(16.2)	(14.5)	(10.6)	(5.9)	(3.9)	(100.0)				
平均	4.30	4.81	4.99	5.15	5.27	5.37	5.45	5.50	5.57	5.62	5.23	5.07	5.22		
標準偏差	1.45	1.42	1.41	1.40	1.39	1.38	1.36	1.32	1.34	1.35	1.42	1.50	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査