

表 3182 %クレアチニン産生速度(%) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	385	405	737	1,174	1,860	2,422	2,520	2,034	1,125	691	13,353	773	14,126	107.12	22.44
(%)	(2.9)	(3.0)	(5.5)	(8.8)	(13.9)	(18.1)	(18.9)	(15.2)	(8.4)	(5.2)	(100.0)				
慢性腎盂腎炎	18	14	33	44	55	60	68	50	25	12	379	19	398	102.47	22.70
(%)	(4.7)	(3.7)	(8.7)	(11.6)	(14.5)	(15.8)	(17.9)	(13.2)	(6.6)	(3.2)	(100.0)				
急速進行性糸球体腎炎	12	6	16	21	37	46	47	29	11	14	239	11	250	104.06	25.24
(%)	(5.0)	(2.5)	(6.7)	(8.8)	(15.5)	(19.2)	(19.7)	(12.1)	(4.6)	(5.9)	(100.0)				
妊娠腎/妊娠中毒症	6	8	14	35	54	60	61	53	30	13	334	14	348	107.39	20.68
(%)	(1.8)	(2.4)	(4.2)	(10.5)	(16.2)	(18.0)	(18.3)	(15.9)	(9.0)	(3.9)	(100.0)				
その他の分類不能の腎炎	4	2	12	16	19	39	49	22	6	9	178	8	186	106.83	20.17
(%)	(2.2)	(1.1)	(6.7)	(9.0)	(10.7)	(21.9)	(27.5)	(12.4)	(3.4)	(5.1)	(100.0)				
多発性嚢胞腎	28	38	86	167	220	241	213	146	74	28	1,241	89	1,330	102.74	20.24
(%)	(2.3)	(3.1)	(6.9)	(13.5)	(17.7)	(19.4)	(17.2)	(11.8)	(6.0)	(2.3)	(100.0)				
腎硬化症	99	79	142	233	335	364	358	254	174	108	2,146	106	2,252	103.79	24.10
(%)	(4.6)	(3.7)	(6.6)	(10.9)	(15.6)	(17.0)	(16.7)	(11.8)	(8.1)	(5.0)	(100.0)				
悪性高血圧	16	8	11	33	39	45	38	32	23	14	259	11	270	104.03	26.29
(%)	(6.2)	(3.1)	(4.2)	(12.7)	(15.1)	(17.4)	(14.7)	(12.4)	(8.9)	(5.4)	(100.0)				
糖尿病性腎症	840	653	1,080	1,654	1,791	1,704	1,240	673	328	214	10,177	668	10,845	93.87	23.86
(%)	(8.3)	(6.4)	(10.6)	(16.3)	(17.6)	(16.7)	(12.2)	(6.6)	(3.2)	(2.1)	(100.0)				
SLE腎炎	20	18	37	45	51	55	44	28	12	7	317	20	337	96.38	22.87
(%)	(6.3)	(5.7)	(11.7)	(14.2)	(16.1)	(17.4)	(13.9)	(8.8)	(3.8)	(2.2)	(100.0)				
アミロイド腎	4	4	5	2	4	10	10	2		1	42	1	43	96.31	27.76
(%)	(9.5)	(9.5)	(11.9)	(4.8)	(9.5)	(23.8)	(23.8)	(4.8)		(2.4)	(100.0)				
痛風腎	2	2	10	13	15	25	25	19	14	6	131	11	142	108.28	21.50
(%)	(1.5)	(1.5)	(7.6)	(9.9)	(11.5)	(19.1)	(19.1)	(14.5)	(10.7)	(4.6)	(100.0)				
先天性代謝異常による腎不全	1	1	1	3	10	5	8	3	2	1	35	6	41	103.35	20.08
(%)	(2.9)	(2.9)	(2.9)	(8.6)	(28.6)	(14.3)	(22.9)	(8.6)	(5.7)	(2.9)	(100.0)				
腎・尿路結核		2	3	1	8	3	1		1		19	1	20	93.65	17.42
(%)		(10.5)	(15.8)	(5.3)	(42.1)	(15.8)	(5.3)		(5.3)		(100.0)				
腎・尿路結石	3	2	4	8	11	11	9	8	5	2	63	1	64	101.93	25.10
(%)	(4.8)	(3.2)	(6.3)	(12.7)	(17.5)	(17.5)	(14.3)	(12.7)	(7.9)	(3.2)	(100.0)				
腎・尿路腫瘍	3	2	3	7	6	10	10	7	6	5	59	3	62	107.03	26.99
(%)	(5.1)	(3.4)	(5.1)	(11.9)	(10.2)	(16.9)	(16.9)	(11.9)	(10.2)	(8.5)	(100.0)				
閉塞性尿路障害		5	2	11	8	14	13	6	2	3	64	5	69	103.33	20.88
(%)		(7.8)	(3.1)	(17.2)	(12.5)	(21.9)	(20.3)	(9.4)	(3.1)	(4.7)	(100.0)				
骨髄腫	3		1		3	3		1			11		11	83.91	30.67
(%)	(27.3)		(9.1)		(27.3)	(27.3)		(9.1)			(100.0)				
腎形成不全	2	3	3	6	14	15	27	14	5	2	91	4	95	107.11	19.85
(%)	(2.2)	(3.3)	(3.3)	(6.6)	(15.4)	(16.5)	(29.7)	(15.4)	(5.5)	(2.2)	(100.0)				
不明	139	108	209	285	385	496	479	327	184	125	2,737	217	2,954	102.92	24.50
(%)	(5.1)	(3.9)	(7.6)	(10.4)	(14.1)	(18.1)	(17.5)	(11.9)	(6.7)	(4.6)	(100.0)				
再導入	5	9	27	33	57	77	70	50	26	14	368	17	385	105.94	20.25
(%)	(1.4)	(2.4)	(7.3)	(9.0)	(15.5)	(20.9)	(19.0)	(13.6)	(7.1)	(3.8)	(100.0)				
その他	44	35	42	77	118	129	109	85	39	39	717	26	743	101.85	25.65
(%)	(6.1)	(4.9)	(5.9)	(10.7)	(16.5)	(18.0)	(15.2)	(11.9)	(5.4)	(5.4)	(100.0)				
合計	1,634	1,404	2,478	3,868	5,100	5,834	5,399	3,843	2,092	1,308	32,960	2,011	34,971	101.92	23.87
(%)	(5.0)	(4.3)	(7.5)	(11.7)	(15.5)	(17.7)	(16.4)	(11.7)	(6.3)	(4.0)	(100.0)				
記載なし															
(%)															
総計	1,634	1,404	2,478	3,868	5,100	5,834	5,399	3,843	2,092	1,308	32,960	2,011	34,971	101.92	23.87
(%)	(5.0)	(4.3)	(7.5)	(11.7)	(15.5)	(17.7)	(16.4)	(11.7)	(6.3)	(4.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査