

表 3190 %クレアチニン産生速度(%) 透析前リン濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
3未満	206	74	97	129	125	108	82	53	34	16	924	44	968	84.71	30.58
(%)	(22.3)	(8.0)	(10.5)	(14.0)	(13.5)	(11.7)	(8.9)	(5.7)	(3.7)	(1.7)	(100.0)				
3.0～	154	96	153	181	190	187	149	94	32	21	1,257	57	1,314	90.39	26.14
(%)	(12.3)	(7.6)	(12.2)	(14.4)	(15.1)	(14.9)	(11.9)	(7.5)	(2.5)	(1.7)	(100.0)				
3.5～	216	142	226	321	357	384	307	205	88	69	2,315	107	2,422	95.38	26.03
(%)	(9.3)	(6.1)	(9.8)	(13.9)	(15.4)	(16.6)	(13.3)	(8.9)	(3.8)	(3.0)	(100.0)				
4.0～	263	208	330	497	628	647	543	404	209	99	3,828	134	3,962	98.45	24.24
(%)	(6.9)	(5.4)	(8.6)	(13.0)	(16.4)	(16.9)	(14.2)	(10.6)	(5.5)	(2.6)	(100.0)				
4.5～	232	211	362	584	759	851	787	542	303	175	4,806	185	4,991	101.68	23.43
(%)	(4.8)	(4.4)	(7.5)	(12.2)	(15.8)	(17.7)	(16.4)	(11.3)	(6.3)	(3.6)	(100.0)				
5.0～	193	234	379	574	834	942	925	647	332	205	5,265	195	5,460	103.05	22.63
(%)	(3.7)	(4.4)	(7.2)	(10.9)	(15.8)	(17.9)	(17.6)	(12.3)	(6.3)	(3.9)	(100.0)				
5.5～	115	162	301	466	687	813	835	600	334	201	4,514	154	4,668	105.21	21.98
(%)	(2.5)	(3.6)	(6.7)	(10.3)	(15.2)	(18.0)	(18.5)	(13.3)	(7.4)	(4.5)	(100.0)				
6.0～	34	28	45	88	124	150	134	107	68	37	815	29	844	104.78	23.44
(%)	(4.2)	(3.4)	(5.5)	(10.8)	(15.2)	(18.4)	(16.4)	(13.1)	(8.3)	(4.5)	(100.0)				
6.1～	72	83	187	301	411	524	488	365	204	142	2,777	107	2,884	105.48	21.94
(%)	(2.6)	(3.0)	(6.7)	(10.8)	(14.8)	(18.9)	(17.6)	(13.1)	(7.3)	(5.1)	(100.0)				
6.5～	54	59	153	296	379	440	403	309	175	124	2,392	82	2,474	105.62	21.98
(%)	(2.3)	(2.5)	(6.4)	(12.4)	(15.8)	(18.4)	(16.8)	(12.9)	(7.3)	(5.2)	(100.0)				
7.0～	27	38	103	164	226	295	281	187	133	83	1,537	50	1,587	106.62	21.69
(%)	(1.8)	(2.5)	(6.7)	(10.7)	(14.7)	(19.2)	(18.3)	(12.2)	(8.7)	(5.4)	(100.0)				
7.5～	30	28	57	86	137	181	159	128	79	65	950	28	978	107.06	23.70
(%)	(3.2)	(2.9)	(6.0)	(9.1)	(14.4)	(19.1)	(16.7)	(13.5)	(8.3)	(6.8)	(100.0)				
8.0～	15	15	34	61	81	118	111	70	36	23	564	14	578	105.17	20.93
(%)	(2.7)	(2.7)	(6.0)	(10.8)	(14.4)	(20.9)	(19.7)	(12.4)	(6.4)	(4.1)	(100.0)				
8.5～	2	11	16	38	56	73	66	50	21	13	346	18	364	106.90	20.54
(%)	(0.6)	(3.2)	(4.6)	(11.0)	(16.2)	(21.1)	(19.1)	(14.5)	(6.1)	(3.8)	(100.0)				
9.0～	14	13	31	77	86	106	111	72	39	33	582	19	601	106.26	22.50
(%)	(2.4)	(2.2)	(5.3)	(13.2)	(14.8)	(18.2)	(19.1)	(12.4)	(6.7)	(5.7)	(100.0)				
合計	1,627	1,402	2,474	3,863	5,080	5,819	5,381	3,833	2,087	1,306	32,872	1,223	34,095	101.92	23.87
(%)	(4.9)	(4.3)	(7.5)	(11.8)	(15.5)	(17.7)	(16.4)	(11.7)	(6.3)	(4.0)	(100.0)				
記載なし	7	2	4	5	20	15	18	10	5	2	88	788	876	101.42	24.52
(%)	(8.0)	(2.3)	(4.5)	(5.7)	(22.7)	(17.0)	(20.5)	(11.4)	(5.7)	(2.3)	(100.0)				
総計	1,634	1,404	2,478	3,868	5,100	5,834	5,399	3,843	2,092	1,308	32,960	2,011	34,971	101.92	23.87
(%)	(5.0)	(4.3)	(7.5)	(11.7)	(15.5)	(17.7)	(16.4)	(11.7)	(6.3)	(4.0)	(100.0)				
平均	4.50	4.94	5.13	5.28	5.36	5.48	5.53	5.55	5.62	5.73	5.38	5.26	5.37		
標準偏差	1.47	1.37	1.40	1.45	1.39	1.41	1.38	1.37	1.32	1.37	1.42	1.44	1.42		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査