

表 3196 %クレアチニン産生速度(%) Non-HDL-C濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

Non-HDL-C濃度	60未満	60～	70～	80～	90～	100～	110～	120～	130～	140～	合計	記載なし	総計	平均	標準偏差
50未満	45	43	41	50	61	85	55	40	15	12	447	16	463	94.02	26.14
(%)	(10.1)	(9.6)	(9.2)	(11.2)	(13.6)	(19.0)	(12.3)	(8.9)	(3.4)	(2.7)	(100.0)				
50～	128	85	189	288	336	338	324	227	95	71	2,081	54	2,135	99.47	24.33
(%)	(6.2)	(4.1)	(9.1)	(13.8)	(16.1)	(16.2)	(15.6)	(10.9)	(4.6)	(3.4)	(100.0)				
70～	238	209	343	554	693	796	678	442	265	192	4,410	123	4,533	100.83	23.93
(%)	(5.4)	(4.7)	(7.8)	(12.6)	(15.7)	(18.0)	(15.4)	(10.0)	(6.0)	(4.4)	(100.0)				
90～	233	210	384	651	854	883	935	618	372	213	5,353	166	5,519	102.65	23.37
(%)	(4.4)	(3.9)	(7.2)	(12.2)	(16.0)	(16.5)	(17.5)	(11.5)	(6.9)	(4.0)	(100.0)				
110～	179	157	315	493	681	781	739	588	319	182	4,434	126	4,560	103.62	23.23
(%)	(4.0)	(3.5)	(7.1)	(11.1)	(15.4)	(17.6)	(16.7)	(13.3)	(7.2)	(4.1)	(100.0)				
130～	96	91	175	272	398	498	506	344	197	115	2,692	74	2,766	104.79	23.09
(%)	(3.6)	(3.4)	(6.5)	(10.1)	(14.8)	(18.5)	(18.8)	(12.8)	(7.3)	(4.3)	(100.0)				
150～	38	45	90	154	202	277	243	170	102	58	1,379	34	1,413	104.60	22.21
(%)	(2.8)	(3.3)	(6.5)	(11.2)	(14.6)	(20.1)	(17.6)	(12.3)	(7.4)	(4.2)	(100.0)				
170～	21	25	34	53	79	100	112	93	44	20	581	14	595	105.18	23.25
(%)	(3.6)	(4.3)	(5.9)	(9.1)	(13.6)	(17.2)	(19.3)	(16.0)	(7.6)	(3.4)	(100.0)				
190～	15	8	12	15	34	35	41	32	7	11	210	6	216	102.97	24.07
(%)	(7.1)	(3.8)	(5.7)	(7.1)	(16.2)	(16.7)	(19.5)	(15.2)	(3.3)	(5.2)	(100.0)				
210～	6	4	9	12	21	20	16	12	9	4	113	2	115	101.43	23.15
(%)	(5.3)	(3.5)	(8.0)	(10.6)	(18.6)	(17.7)	(14.2)	(10.6)	(8.0)	(3.5)	(100.0)				
合計	999	877	1,592	2,542	3,359	3,813	3,649	2,566	1,425	878	21,700	615	22,315	102.45	23.59
(%)	(4.6)	(4.0)	(7.3)	(11.7)	(15.5)	(17.6)	(16.8)	(11.8)	(6.6)	(4.0)	(100.0)				
記載なし	635	527	886	1,326	1,741	2,021	1,750	1,277	667	430	11,260	1,396	12,656	100.91	24.38
(%)	(5.6)	(4.7)	(7.9)	(11.8)	(15.5)	(17.9)	(15.5)	(11.3)	(5.9)	(3.8)	(100.0)				
総計	1,634	1,404	2,478	3,868	5,100	5,834	5,399	3,843	2,092	1,308	32,960	2,011	34,971	101.92	23.87
(%)	(5.0)	(4.3)	(7.5)	(11.7)	(15.5)	(17.7)	(16.4)	(11.7)	(6.3)	(4.0)	(100.0)				
平均	100.86	102.52	104.02	103.90	106.20	107.86	109.12	110.20	110.63	107.98	106.99	105.96	106.97		
標準偏差	34.90	34.65	33.14	32.31	32.78	33.29	33.43	33.39	31.66	32.26	33.22	32.06	33.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査