

表 3213 糖尿病の有無 PTH値(pg/mL)別（透析患者全体）PTH測定法:intact-PTH

PTH値	糖尿病なし	糖尿病あり	合計	不明	記載なし	総計
20未満	5,414	5,462	10,876	997		11,873
(%)	(49.8)	(50.2)	(100.0)			
20～	7,343	8,187	15,530	1,416		16,946
(%)	(47.3)	(52.7)	(100.0)			
40～	7,884	8,604	16,488	1,528		18,016
(%)	(47.8)	(52.2)	(100.0)			
60～	8,587	9,052	17,639	1,591		19,230
(%)	(48.7)	(51.3)	(100.0)			
80～	8,799	9,134	17,933	1,680		19,613
(%)	(49.1)	(50.9)	(100.0)			
100～	8,503	8,922	17,425	1,733		19,158
(%)	(48.8)	(51.2)	(100.0)			
120～	8,322	8,250	16,572	1,650		18,222
(%)	(50.2)	(49.8)	(100.0)			
140～	7,441	7,834	15,275	1,532		16,807
(%)	(48.7)	(51.3)	(100.0)			
160～	6,850	6,973	13,823	1,350		15,173
(%)	(49.6)	(50.4)	(100.0)			
180～	15,918	15,478	31,396	3,117		34,513
(%)	(50.7)	(49.3)	(100.0)			
240～	199	207	406	38		444
(%)	(49.0)	(51.0)	(100.0)			
241～	9,347	8,626	17,973	1,866		19,839
(%)	(52.0)	(48.0)	(100.0)			
300～	5,519	4,922	10,441	1,085		11,526
(%)	(52.9)	(47.1)	(100.0)			
360～	3,284	2,723	6,007	677		6,684
(%)	(54.7)	(45.3)	(100.0)			
420～	1,970	1,540	3,510	415		3,925
(%)	(56.1)	(43.9)	(100.0)			
480～	509	363	872	92		964
(%)	(58.4)	(41.6)	(100.0)			
500～	810	581	1,391	173		1,564
(%)	(58.2)	(41.8)	(100.0)			
540～	891	637	1,528	186		1,714
(%)	(58.3)	(41.7)	(100.0)			
600～	1,599	911	2,510	289		2,799
(%)	(63.7)	(36.3)	(100.0)			
800～	1,382	548	1,930	232		2,162
(%)	(71.6)	(28.4)	(100.0)			
合計	110,571	108,954	219,525	21,647		241,172
(%)	(50.4)	(49.6)	(100.0)			
記載なし	2,153	3,160	5,313	1,071		6,384
(%)	(40.5)	(59.5)	(100.0)			
総計	112,724	112,114	224,838	22,718		247,556
(%)	(50.1)	(49.9)	(100.0)			
平均	181.39	161.04	171.29	181.11		172.17
標準偏差	182.50	138.77	162.60	176.79		163.94

数値下のかっこ内は行方向の合計に対する%です。

*糖尿病ありとは、原疾患が糖尿病性腎症または、糖尿病の既往が有りの患者。

糖尿病なしとは、原疾患が糖尿病性腎症以外で且つ糖尿病の既往がなしの患者。

記載なしとは、原疾患と糖尿病の既往いずれも記載のなかった患者。

不明とは、上記以外。

(C)Japanese Society for Dialysis Therapy

2014年調査