

表 3224 透析前アルブミン濃度(g/dL) 併用の状況別 (透析患者全体)

併用の状況	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	170	914	3,747	14,349	63,843	133,228	50,050	2,891	201	269,393	31,401	300,794	3.60	0.44
(%)	(0.1)	(0.3)	(1.4)	(5.3)	(23.7)	(49.5)	(18.6)	(1.1)	(0.1)	(100.0)				
PD単独	12	65	254	785	1,591	1,519	366	18	1	4,611	2,577	7,188	3.28	0.54
(%)	(0.3)	(1.4)	(5.5)	(17.0)	(34.5)	(32.9)	(7.9)	(0.4)	(0.0)	(100.0)				
HD(F)単独.PDカテあり		2	15	35	77	71	16	1		217	52	269	3.28	0.53
(%)		(0.9)	(6.9)	(16.1)	(35.5)	(32.7)	(7.4)	(0.5)		(100.0)				
PD.HD併用週1 HD(F)	2	3	42	237	506	325	56	3	1	1,175	369	1,544	3.25	0.45
(%)	(0.2)	(0.3)	(3.6)	(20.2)	(43.1)	(27.7)	(4.8)	(0.3)	(0.1)	(100.0)				
PD.HD併用週2 HD(F)		1	1	8	54	60	12			136	41	177	3.45	0.40
(%)		(0.7)	(0.7)	(5.9)	(39.7)	(44.1)	(8.8)			(100.0)				
PD.HD併用週3 HD(F)		2		8	9	11	3	1		34	6	40	3.28	0.65
(%)		(5.9)		(23.5)	(26.5)	(32.4)	(8.8)	(2.9)		(100.0)				
PD.HD併用週4 HD(F)				1						1		1	2.90	
(%)				(100.0)						(100.0)				
PD.HD併用その他	1	2	1	11	25	16	2			58	15	73	3.19	0.55
(%)	(1.7)	(3.4)	(1.7)	(19.0)	(43.1)	(27.6)	(3.4)			(100.0)				
合計	185	989	4,060	15,434	66,105	135,230	50,505	2,914	203	275,625	34,461	310,086	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				
不明			1	2	3	5	3			14	8	22	3.46	0.54
(%)			(7.1)	(14.3)	(21.4)	(35.7)	(21.4)			(100.0)				
記載なし														
(%)														
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査