

表 3228 透析前アルブミン濃度(g/dL) 原疾患別（透析患者全体）

原疾患	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	46	196	934	4,055	19,733	43,438	16,702	1,009	68	86,181	10,789	96,970	3.63	0.42
(%)	(0.1)	(0.2)	(1.1)	(4.7)	(22.9)	(50.4)	(19.4)	(1.2)	(0.1)	(100.0)				
慢性腎盂腎炎	1	16	42	140	659	1,350	492	32	6	2,738	304	3,042	3.60	0.46
(%)	(0.0)	(0.6)	(1.5)	(5.1)	(24.1)	(49.3)	(18.0)	(1.2)	(0.2)	(100.0)				
急速進行性糸球体腎炎		16	51	194	645	979	289	20	1	2,195	299	2,494	3.49	0.48
(%)		(0.7)	(2.3)	(8.8)	(29.4)	(44.6)	(13.2)	(0.9)	(0.0)	(100.0)				
妊娠腎/妊娠中毒症			5	56	313	812	272	11	1	1,470	144	1,614	3.65	0.36
(%)			(0.3)	(3.8)	(21.3)	(55.2)	(18.5)	(0.7)	(0.1)	(100.0)				
その他の分類不能の腎炎		5	21	52	238	595	289	34	2	1,236	137	1,373	3.68	0.46
(%)		(0.4)	(1.7)	(4.2)	(19.3)	(48.1)	(23.4)	(2.8)	(0.2)	(100.0)				
多発性嚢胞腎	2	37	95	449	2,364	5,044	1,801	83	5	9,880	1,126	11,006	3.61	0.41
(%)	(0.0)	(0.4)	(1.0)	(4.5)	(23.9)	(51.1)	(18.2)	(0.8)	(0.1)	(100.0)				
腎硬化症	12	80	424	1,641	6,934	12,216	3,792	229	23	25,351	2,947	28,298	3.55	0.44
(%)	(0.0)	(0.3)	(1.7)	(6.5)	(27.4)	(48.2)	(15.0)	(0.9)	(0.1)	(100.0)				
悪性高血圧	2	4	17	103	435	1,130	533	45	1	2,270	303	2,573	3.68	0.43
(%)	(0.1)	(0.2)	(0.7)	(4.5)	(19.2)	(49.8)	(23.5)	(2.0)	(0.0)	(100.0)				
糖尿病性腎症	85	430	1,703	6,170	25,009	51,203	19,460	981	61	105,102	12,979	118,081	3.59	0.45
(%)	(0.1)	(0.4)	(1.6)	(5.9)	(23.8)	(48.7)	(18.5)	(0.9)	(0.1)	(100.0)				
SLE腎炎	3	8	33	119	601	924	277	16		1,981	255	2,236	3.53	0.44
(%)	(0.2)	(0.4)	(1.7)	(6.0)	(30.3)	(46.6)	(14.0)	(0.8)		(100.0)				
アミロイド腎	1	3	16	49	105	152	58			384	67	451	3.43	0.54
(%)	(0.3)	(0.8)	(4.2)	(12.8)	(27.3)	(39.6)	(15.1)			(100.0)				
痛風腎			7	35	222	531	197	7		999	114	1,113	3.65	0.38
(%)			(0.7)	(3.5)	(22.2)	(53.2)	(19.7)	(0.7)		(100.0)				
先天性代謝異常による腎不全		1	1	16	44	90	73	7	3	235	37	272	3.74	0.52
(%)		(0.4)	(0.4)	(6.8)	(18.7)	(38.3)	(31.1)	(3.0)	(1.3)	(100.0)				
腎・尿路結核		1	2	19	51	78	26			177	29	206	3.50	0.45
(%)		(0.6)	(1.1)	(10.7)	(28.8)	(44.1)	(14.7)			(100.0)				
腎・尿路結石		1	9	38	130	249	92	2	1	522	59	581	3.57	0.45
(%)		(0.2)	(1.7)	(7.3)	(24.9)	(47.7)	(17.6)	(0.4)	(0.2)	(100.0)				
腎・尿路腫瘍		6	14	44	232	379	102	6		783	103	886	3.53	0.45
(%)		(0.8)	(1.8)	(5.6)	(29.6)	(48.4)	(13.0)	(0.8)		(100.0)				
閉塞性尿路障害		3	12	35	146	306	124	8	1	635	88	723	3.61	0.47
(%)		(0.5)	(1.9)	(5.5)	(23.0)	(48.2)	(19.5)	(1.3)	(0.2)	(100.0)				
骨髄腫	2	2	8	30	56	91	39	4		232	44	276	3.47	0.57
(%)	(0.9)	(0.9)	(3.4)	(12.9)	(24.1)	(39.2)	(16.8)	(1.7)		(100.0)				
腎形成不全		1	2	15	85	266	182	22	1	574	75	649	3.79	0.42
(%)		(0.2)	(0.3)	(2.6)	(14.8)	(46.3)	(31.7)	(3.8)	(0.2)	(100.0)				
不明	23	128	504	1,623	6,102	11,403	4,037	264	21	24,105	3,439	27,544	3.56	0.47
(%)	(0.1)	(0.5)	(2.1)	(6.7)	(25.3)	(47.3)	(16.7)	(1.1)	(0.1)	(100.0)				
再導入	2	1	19	78	393	960	455	29	5	1,942	221	2,163	3.68	0.43
(%)	(0.1)	(0.1)	(1.0)	(4.0)	(20.2)	(49.4)	(23.4)	(1.5)	(0.3)	(100.0)				
その他	6	50	142	475	1,611	3,039	1,216	105	3	6,647	910	7,557	3.56	0.49
(%)	(0.1)	(0.8)	(2.1)	(7.1)	(24.2)	(45.7)	(18.3)	(1.6)	(0.0)	(100.0)				
合計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				
記載なし														
(%)														
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査