

表 3229 透析前アルブミン濃度(g/dL) 医療機関別（透析患者全体）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	6	22	100	157	301	310	61	5	1	963	294	1,257	3.21	0.60
(%)	(0.6)	(2.3)	(10.4)	(16.3)	(31.3)	(32.2)	(6.3)	(0.5)	(0.1)	(100.0)				
私立大学	13	56	122	325	674	960	354	26	2	2,532	985	3,517	3.39	0.59
(%)	(0.5)	(2.2)	(4.8)	(12.8)	(26.6)	(37.9)	(14.0)	(1.0)	(0.1)	(100.0)				
国立	1	16	28	102	198	266	78	8	1	698	171	869	3.38	0.58
(%)	(0.1)	(2.3)	(4.0)	(14.6)	(28.4)	(38.1)	(11.2)	(1.1)	(0.1)	(100.0)				
県市町村立	22	134	452	1,310	4,391	7,027	2,204	149	13	15,702	3,483	19,185	3.49	0.50
(%)	(0.1)	(0.9)	(2.9)	(8.3)	(28.0)	(44.8)	(14.0)	(0.9)	(0.1)	(100.0)				
社会保険	5	32	105	282	1,049	1,492	462	21	2	3,450	422	3,872	3.47	0.49
(%)	(0.1)	(0.9)	(3.0)	(8.2)	(30.4)	(43.2)	(13.4)	(0.6)	(0.1)	(100.0)				
厚生連	11	55	171	664	2,343	3,491	1,250	89	4	8,078	1,186	9,264	3.52	0.48
(%)	(0.1)	(0.7)	(2.1)	(8.2)	(29.0)	(43.2)	(15.5)	(1.1)	(0.0)	(100.0)				
その他公的	11	60	233	776	2,507	4,090	1,299	83	8	9,067	1,922	10,989	3.50	0.49
(%)	(0.1)	(0.7)	(2.6)	(8.6)	(27.6)	(45.1)	(14.3)	(0.9)	(0.1)	(100.0)				
私立総合	12	39	148	496	1,713	2,969	1,014	48	3	6,442	1,112	7,554	3.53	0.48
(%)	(0.2)	(0.6)	(2.3)	(7.7)	(26.6)	(46.1)	(15.7)	(0.7)	(0.0)	(100.0)				
私立	85	441	1,927	6,395	20,885	35,016	11,823	666	69	77,307	11,335	88,642	3.52	0.48
(%)	(0.1)	(0.6)	(2.5)	(8.3)	(27.0)	(45.3)	(15.3)	(0.9)	(0.1)	(100.0)				
私立診療所	19	134	775	4,929	32,047	79,614	31,963	1,819	100	151,400	13,559	164,959	3.67	0.39
(%)	(0.0)	(0.1)	(0.5)	(3.3)	(21.2)	(52.6)	(21.1)	(1.2)	(0.1)	(100.0)				
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査