

表 3230 透析前アルブミン濃度(g/dL) 週透析回数(回/週)別(透析患者全体)

週透析回数	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
1回	1	13	54	289	624	519	160	19	1	1,680	160	1,840	3.34	0.51
(%)	(0.1)	(0.8)	(3.2)	(17.2)	(37.1)	(30.9)	(9.5)	(1.1)	(0.1)	(100.0)				
2回	11	54	160	454	1,363	2,491	1,188	103	6	5,830	362	6,192	3.56	0.53
(%)	(0.2)	(0.9)	(2.7)	(7.8)	(23.4)	(42.7)	(20.4)	(1.8)	(0.1)	(100.0)				
3回	156	834	3,490	13,607	61,331	128,590	47,923	2,695	193	258,819	11,820	270,639	3.60	0.44
(%)	(0.1)	(0.3)	(1.3)	(5.3)	(23.7)	(49.7)	(18.5)	(1.0)	(0.1)	(100.0)				
4回	1	5	23	49	208	435	210	22		953	41	994	3.63	0.50
(%)	(0.1)	(0.5)	(2.4)	(5.1)	(21.8)	(45.6)	(22.0)	(2.3)		(100.0)				
5回			1	4	11	29	28	7	1	81	8	89	3.82	0.54
(%)			(1.2)	(4.9)	(13.6)	(35.8)	(34.6)	(8.6)	(1.2)	(100.0)				
6回		2	2	2	5	26	25	8		70	11	81	3.81	0.63
(%)		(2.9)	(2.9)	(2.9)	(7.1)	(37.1)	(35.7)	(11.4)		(100.0)				
7回					4	6	13	4		27	3	30	4.01	0.41
(%)					(14.8)	(22.2)	(48.1)	(14.8)		(100.0)				
合計	169	908	3,730	14,405	63,546	132,096	49,547	2,858	201	267,460	12,405	279,865	3.60	0.44
(%)	(0.1)	(0.3)	(1.4)	(5.4)	(23.8)	(49.4)	(18.5)	(1.1)	(0.1)	(100.0)				
記載なし	16	81	331	1,031	2,562	3,139	961	56	2	8,179	22,064	30,243	3.39	0.52
(%)	(0.2)	(1.0)	(4.0)	(12.6)	(31.3)	(38.4)	(11.7)	(0.7)	(0.0)	(100.0)				
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				
平均	2.93	2.92	2.94	2.93	2.96	2.98	2.98	2.98	2.97	2.97	2.95	2.97		
標準偏差	0.30	0.37	0.33	0.34	0.26	0.20	0.23	0.36	0.26	0.23	0.31	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査