

表 3233 透析前アルブミン濃度(g/dL) 透析前リン濃度(mg/dL)別 (透析患者全体)

透析前リン濃度	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
3未満	64	285	1,001	1,995	3,279	3,149	836	50	10	10,669	217	10,886	3.19	0.61
(%)	(0.6)	(2.7)	(9.4)	(18.7)	(30.7)	(29.5)	(7.8)	(0.5)	(0.1)	(100.0)				
3.0～	30	111	488	1,617	4,205	5,302	1,485	85	10	13,333	249	13,582	3.41	0.51
(%)	(0.2)	(0.8)	(3.7)	(12.1)	(31.5)	(39.8)	(11.1)	(0.6)	(0.1)	(100.0)				
3.5～	19	117	535	2,045	6,737	10,400	3,283	168	11	23,315	367	23,682	3.50	0.47
(%)	(0.1)	(0.5)	(2.3)	(8.8)	(28.9)	(44.6)	(14.1)	(0.7)	(0.0)	(100.0)				
4.0～	12	118	494	2,255	9,175	16,684	5,683	296	23	34,740	542	35,282	3.57	0.44
(%)	(0.0)	(0.3)	(1.4)	(6.5)	(26.4)	(48.0)	(16.4)	(0.9)	(0.1)	(100.0)				
4.5～	13	100	419	1,984	10,210	20,924	7,369	421	22	41,462	597	42,059	3.61	0.42
(%)	(0.0)	(0.2)	(1.0)	(4.8)	(24.6)	(50.5)	(17.8)	(1.0)	(0.1)	(100.0)				
5.0～	10	68	328	1,738	9,759	21,581	8,114	439	30	42,067	651	42,718	3.64	0.40
(%)	(0.0)	(0.2)	(0.8)	(4.1)	(23.2)	(51.3)	(19.3)	(1.0)	(0.1)	(100.0)				
5.5～	15	50	246	1,285	7,869	18,876	7,424	396	27	36,188	517	36,705	3.66	0.40
(%)	(0.0)	(0.1)	(0.7)	(3.6)	(21.7)	(52.2)	(20.5)	(1.1)	(0.1)	(100.0)				
6.0～	1	8	37	212	1,342	3,209	1,269	85	6	6,169	92	6,261	3.66	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.4)	(21.8)	(52.0)	(20.6)	(1.4)	(0.1)	(100.0)				
6.1～	6	29	113	725	4,313	10,877	4,469	245	15	20,792	317	21,109	3.67	0.39
(%)	(0.0)	(0.1)	(0.5)	(3.5)	(20.7)	(52.3)	(21.5)	(1.2)	(0.1)	(100.0)				
6.5～	6	26	100	561	3,594	9,430	4,000	239	18	17,974	304	18,278	3.68	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.1)	(20.0)	(52.5)	(22.3)	(1.3)	(0.1)	(100.0)				
7.0～	1	12	66	329	2,133	5,869	2,470	192	11	11,083	194	11,277	3.69	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.0)	(19.2)	(53.0)	(22.3)	(1.7)	(0.1)	(100.0)				
7.5～	2	5	40	204	1,344	3,443	1,544	96	6	6,684	174	6,858	3.69	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.1)	(20.1)	(51.5)	(23.1)	(1.4)	(0.1)	(100.0)				
8.0～		10	31	129	725	2,028	958	70	3	3,954	99	4,053	3.70	0.41
(%)		(0.3)	(0.8)	(3.3)	(18.3)	(51.3)	(24.2)	(1.8)	(0.1)	(100.0)				
8.5～		5	19	80	479	1,312	581	52	1	2,529	43	2,572	3.69	0.40
(%)		(0.2)	(0.8)	(3.2)	(18.9)	(51.9)	(23.0)	(2.1)	(0.0)	(100.0)				
9.0～	1	9	30	112	762	2,000	978	73	9	3,974	103	4,077	3.71	0.41
(%)	(0.0)	(0.2)	(0.8)	(2.8)	(19.2)	(50.3)	(24.6)	(1.8)	(0.2)	(100.0)				
合計	180	953	3,947	15,271	65,926	135,084	50,463	2,907	202	274,933	4,466	279,399	3.60	0.44
(%)	(0.1)	(0.3)	(1.4)	(5.6)	(24.0)	(49.1)	(18.4)	(1.1)	(0.1)	(100.0)				
記載なし	5	36	114	165	182	151	45	7	1	706	30,003	30,709	3.01	0.68
(%)	(0.7)	(5.1)	(16.1)	(23.4)	(25.8)	(21.4)	(6.4)	(1.0)	(0.1)	(100.0)				
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				
平均	3.60	3.90	4.03	4.51	5.02	5.32	5.48	5.60	5.45	5.21	5.28	5.21		
標準偏差	1.73	1.72	1.63	1.49	1.40	1.37	1.39	1.47	1.69	1.42	1.58	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査