

表 3234 透析前アルブミン濃度(g/dL) PTH値(pg/mL)別 (透析患者全体) PTH測定法:intact-PTH

PTH値	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
20未満	18	85	337	1,001	3,135	5,351	1,691	88	18	11,724	149	11,873	3.50	0.49
(%)	(0.2)	(0.7)	(2.9)	(8.5)	(26.7)	(45.6)	(14.4)	(0.8)	(0.2)	(100.0)				
20～	14	90	399	1,273	4,195	7,913	2,696	138	14	16,732	214	16,946	3.54	0.48
(%)	(0.1)	(0.5)	(2.4)	(7.6)	(25.1)	(47.3)	(16.1)	(0.8)	(0.1)	(100.0)				
40～	19	88	357	1,235	4,435	8,450	3,040	173	21	17,818	198	18,016	3.56	0.47
(%)	(0.1)	(0.5)	(2.0)	(6.9)	(24.9)	(47.4)	(17.1)	(1.0)	(0.1)	(100.0)				
60～	7	68	310	1,159	4,753	9,103	3,444	196	14	19,054	176	19,230	3.58	0.45
(%)	(0.0)	(0.4)	(1.6)	(6.1)	(24.9)	(47.8)	(18.1)	(1.0)	(0.1)	(100.0)				
80～	13	62	265	1,075	4,810	9,538	3,459	190	6	19,418	195	19,613	3.59	0.43
(%)	(0.1)	(0.3)	(1.4)	(5.5)	(24.8)	(49.1)	(17.8)	(1.0)	(0.0)	(100.0)				
100～	11	56	241	1,029	4,518	9,331	3,597	178	15	18,976	182	19,158	3.60	0.43
(%)	(0.1)	(0.3)	(1.3)	(5.4)	(23.8)	(49.2)	(19.0)	(0.9)	(0.1)	(100.0)				
120～	5	45	197	930	4,222	9,033	3,419	204	9	18,064	158	18,222	3.62	0.42
(%)	(0.0)	(0.2)	(1.1)	(5.1)	(23.4)	(50.0)	(18.9)	(1.1)	(0.0)	(100.0)				
140～	9	29	154	762	3,940	8,405	3,173	179	8	16,659	148	16,807	3.62	0.41
(%)	(0.1)	(0.2)	(0.9)	(4.6)	(23.7)	(50.5)	(19.0)	(1.1)	(0.0)	(100.0)				
160～	6	19	135	672	3,493	7,635	2,906	157	12	15,035	138	15,173	3.63	0.41
(%)	(0.0)	(0.1)	(0.9)	(4.5)	(23.2)	(50.8)	(19.3)	(1.0)	(0.1)	(100.0)				
180～	9	64	246	1,372	7,776	17,497	6,784	368	22	34,138	375	34,513	3.64	0.40
(%)	(0.0)	(0.2)	(0.7)	(4.0)	(22.8)	(51.3)	(19.9)	(1.1)	(0.1)	(100.0)				
240～	1	4	15	97	233	88	6	444				444	3.64	0.42
(%)	(0.2)		(0.9)	(3.4)	(21.8)	(52.5)	(19.8)	(1.4)		(100.0)				
241～	8	29	156	793	4,426	9,906	3,992	269	20	19,599	240	19,839	3.65	0.41
(%)	(0.0)	(0.1)	(0.8)	(4.0)	(22.6)	(50.5)	(20.4)	(1.4)	(0.1)	(100.0)				
300～	4	15	81	452	2,446	5,917	2,306	148	8	11,377	149	11,526	3.65	0.40
(%)	(0.0)	(0.1)	(0.7)	(4.0)	(21.5)	(52.0)	(20.3)	(1.3)	(0.1)	(100.0)				
360～	4	4	42	239	1,497	3,321	1,383	82	5	6,577	107	6,684	3.66	0.40
(%)	(0.1)	(0.1)	(0.6)	(3.6)	(22.8)	(50.5)	(21.0)	(1.2)	(0.1)	(100.0)				
420～		5	30	157	849	1,960	813	63	5	3,882	43	3,925	3.66	0.41
(%)		(0.1)	(0.8)	(4.0)	(21.9)	(50.5)	(20.9)	(1.6)	(0.1)	(100.0)				
480～	1	2	3	55	208	458	213	12	1	953	11	964	3.64	0.43
(%)	(0.1)	(0.2)	(0.3)	(5.8)	(21.8)	(48.1)	(22.4)	(1.3)	(0.1)	(100.0)				
500～	1	2	18	68	351	779	312	16	1	1,548	16	1,564	3.63	0.41
(%)	(0.1)	(0.1)	(1.2)	(4.4)	(22.7)	(50.3)	(20.2)	(1.0)	(0.1)	(100.0)				
540～		15	60	373	873	335	24	2	1,682	32	1,714	3.66	0.40	
(%)		(0.9)	(3.6)	(22.2)	(51.9)	(19.9)	(1.4)	(0.1)	(100.0)					
600～		1	24	94	642	1,404	546	41	6	2,758	41	2,799	3.65	0.41
(%)		(0.0)	(0.9)	(3.4)	(23.3)	(50.9)	(19.8)	(1.5)	(0.2)	(100.0)				
800～		2	24	103	492	1,044	429	24	2,118	44	2,162	3.63	0.42	
(%)		(0.1)	(1.1)	(4.9)	(23.2)	(49.3)	(20.3)	(1.1)		(100.0)				
合計	130	666	3,038	12,544	56,658	118,151	44,626	2,556	187	238,556	2,616	241,172	3.61	0.43
(%)	(0.1)	(0.3)	(1.3)	(5.3)	(23.8)	(49.5)	(18.7)	(1.1)	(0.1)	(100.0)				
記載なし	16	75	191	515	1,298	1,863	623	34	2	4,617	1,767	6,384	3.42	0.55
(%)	(0.3)	(1.6)	(4.1)	(11.2)	(28.1)	(40.4)	(13.5)	(0.7)	(0.0)	(100.0)				
総計	146	741	3,229	13,059	57,956	120,014	45,249	2,590	189	243,173	4,383	247,556	3.60	0.44
(%)	(0.1)	(0.3)	(1.3)	(5.4)	(23.8)	(49.4)	(18.6)	(1.1)	(0.1)	(100.0)				
平均	119.77	108.17	127.84	146.53	167.36	174.76	180.68	190.79	171.34	171.98	189.83	172.17		
標準偏差	105.74	107.83	150.77	160.75	168.93	161.20	161.90	184.78	155.99	163.46	202.24	163.94		

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査