

表 3239 透析前アルブミン濃度(g/dL) Non-HDL-C濃度(mg/dL)別 (透析患者全体)

Non-HDL-C濃度	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
50未満	17	58	139	354	898	1,421	460	41	3	3,391	19	3,410	3.43	0.57
(%)	(0.5)	(1.7)	(4.1)	(10.4)	(26.5)	(41.9)	(13.6)	(1.2)	(0.1)	(100.0)				
50～	15	113	375	1,236	4,314	7,274	2,491	144	19	15,981	120	16,101	3.52	0.48
(%)	(0.1)	(0.7)	(2.3)	(7.7)	(27.0)	(45.5)	(15.6)	(0.9)	(0.1)	(100.0)				
70～	20	122	546	2,153	9,241	16,929	5,943	311	23	35,288	281	35,569	3.57	0.44
(%)	(0.1)	(0.3)	(1.5)	(6.1)	(26.2)	(48.0)	(16.8)	(0.9)	(0.1)	(100.0)				
90～	14	99	410	2,076	10,105	21,071	7,817	440	19	42,051	327	42,378	3.61	0.42
(%)	(0.0)	(0.2)	(1.0)	(4.9)	(24.0)	(50.1)	(18.6)	(1.0)	(0.0)	(100.0)				
110～	17	55	320	1,489	7,717	17,161	6,827	353	31	33,970	233	34,203	3.64	0.41
(%)	(0.1)	(0.2)	(0.9)	(4.4)	(22.7)	(50.5)	(20.1)	(1.0)	(0.1)	(100.0)				
130～	3	38	154	825	4,430	10,806	4,253	270	21	20,800	163	20,963	3.66	0.40
(%)	(0.0)	(0.2)	(0.7)	(4.0)	(21.3)	(52.0)	(20.4)	(1.3)	(0.1)	(100.0)				
150～	3	17	85	408	2,108	5,384	2,363	126	13	10,507	83	10,590	3.67	0.41
(%)	(0.0)	(0.2)	(0.8)	(3.9)	(20.1)	(51.2)	(22.5)	(1.2)	(0.1)	(100.0)				
170～	3	5	44	185	935	2,362	1,061	73	4	4,672	50	4,722	3.67	0.42
(%)	(0.1)	(0.1)	(0.9)	(4.0)	(20.0)	(50.6)	(22.7)	(1.6)	(0.1)	(100.0)				
190～	4	3	31	94	377	852	402	38		1,801	8	1,809	3.64	0.46
(%)	(0.2)	(0.2)	(1.7)	(5.2)	(20.9)	(47.3)	(22.3)	(2.1)		(100.0)				
210～	5	5	29	64	256	509	221	23	1	1,113	11	1,124	3.59	0.51
(%)	(0.4)	(0.4)	(2.6)	(5.8)	(23.0)	(45.7)	(19.9)	(2.1)	(0.1)	(100.0)				
合計	101	515	2,133	8,884	40,381	83,769	31,838	1,819	134	169,574	1,295	170,869	3.61	0.43
(%)	(0.1)	(0.3)	(1.3)	(5.2)	(23.8)	(49.4)	(18.8)	(1.1)	(0.1)	(100.0)				
記載なし	84	474	1,928	6,552	25,727	51,466	18,670	1,095	69	106,065	33,174	139,239	3.58	0.46
(%)	(0.1)	(0.4)	(1.8)	(6.2)	(24.3)	(48.5)	(17.6)	(1.0)	(0.1)	(100.0)				
総計	185	989	4,061	15,436	66,108	135,235	50,508	2,914	203	275,639	34,469	310,108	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(24.0)	(49.1)	(18.3)	(1.1)	(0.1)	(100.0)				
平均	98.40	88.85	96.98	100.17	104.21	108.11	110.64	112.18	110.05	107.08	107.61	107.08		
標準偏差	53.80	38.01	39.10	34.78	33.07	33.11	33.65	37.13	38.17	33.61	33.84	33.61		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査