

表 3246 透析前アルブミン濃度(g/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	33	154	729	3,045	14,727	32,946	12,866	796	58	65,354	3,223	68,577	3.63	0.42
(%)	(0.1)	(0.2)	(1.1)	(4.7)	(22.5)	(50.4)	(19.7)	(1.2)	(0.1)	(100.0)				
慢性腎盂腎炎		11	33	109	522	1,005	370	24	4	2,078	87	2,165	3.59	0.46
(%)		(0.5)	(1.6)	(5.2)	(25.1)	(48.4)	(17.8)	(1.2)	(0.2)	(100.0)				
急速進行性糸球体腎炎		12	41	154	537	775	217	10	1	1,747	87	1,834	3.48	0.47
(%)		(0.7)	(2.3)	(8.8)	(30.7)	(44.4)	(12.4)	(0.6)	(0.1)	(100.0)				
妊娠腎/妊娠中毒症			5	42	207	548	199	7	1	1,009	55	1,064	3.66	0.37
(%)			(0.5)	(4.2)	(20.5)	(54.3)	(19.7)	(0.7)	(0.1)	(100.0)				
その他の分類不能の腎炎		3	16	36	173	440	212	26	1	907	42	949	3.68	0.45
(%)		(0.3)	(1.8)	(4.0)	(19.1)	(48.5)	(23.4)	(2.9)	(0.1)	(100.0)				
多発性嚢胞腎	2	27	72	345	1,885	3,993	1,453	65	4	7,846	352	8,198	3.62	0.41
(%)	(0.0)	(0.3)	(0.9)	(4.4)	(24.0)	(50.9)	(18.5)	(0.8)	(0.1)	(100.0)				
腎硬化症	11	61	332	1,295	5,600	10,011	3,074	185	22	20,591	869	21,460	3.55	0.44
(%)	(0.1)	(0.3)	(1.6)	(6.3)	(27.2)	(48.6)	(14.9)	(0.9)	(0.1)	(100.0)				
悪性高血圧	1	4	16	80	345	873	423	37	1	1,780	101	1,881	3.68	0.44
(%)	(0.1)	(0.2)	(0.9)	(4.5)	(19.4)	(49.0)	(23.8)	(2.1)	(0.1)	(100.0)				
糖尿病性腎症	73	340	1,321	4,974	20,378	42,180	16,299	837	55	86,457	4,050	90,507	3.60	0.45
(%)	(0.1)	(0.4)	(1.5)	(5.8)	(23.6)	(48.8)	(18.9)	(1.0)	(0.1)	(100.0)				
SLE腎炎	2	6	25	99	459	679	203	14		1,487	63	1,550	3.52	0.44
(%)	(0.1)	(0.4)	(1.7)	(6.7)	(30.9)	(45.7)	(13.7)	(0.9)		(100.0)				
アミロイド腎	1	2	13	39	86	119	39			299	26	325	3.41	0.53
(%)	(0.3)	(0.7)	(4.3)	(13.0)	(28.8)	(39.8)	(13.0)			(100.0)				
痛風腎			6	30	163	420	165	6		790	38	828	3.66	0.38
(%)			(0.8)	(3.8)	(20.6)	(53.2)	(20.9)	(0.8)		(100.0)				
先天性代謝異常による腎不全		1	1	11	27	66	55	5	3	169	8	177	3.76	0.53
(%)		(0.6)	(0.6)	(6.5)	(16.0)	(39.1)	(32.5)	(3.0)	(1.8)	(100.0)				
腎・尿路結核		1	2	18	44	61	22			148	10	158	3.48	0.47
(%)		(0.7)	(1.4)	(12.2)	(29.7)	(41.2)	(14.9)			(100.0)				
腎・尿路結石		1	7	30	101	204	69	1	1	414	16	430	3.57	0.45
(%)		(0.2)	(1.7)	(7.2)	(24.4)	(49.3)	(16.7)	(0.2)	(0.2)	(100.0)				
腎・尿路腫瘍		6	14	35	196	312	90	4		657	29	686	3.52	0.46
(%)		(0.9)	(2.1)	(5.3)	(29.8)	(47.5)	(13.7)	(0.6)		(100.0)				
閉塞性尿路障害		2	10	28	110	240	97	7		494	23	517	3.61	0.47
(%)		(0.4)	(2.0)	(5.7)	(22.3)	(48.6)	(19.6)	(1.4)		(100.0)				
骨髄腫	2	2	6	26	46	66	25	3		176	14	190	3.42	0.59
(%)	(1.1)	(1.1)	(3.4)	(14.8)	(26.1)	(37.5)	(14.2)	(1.7)		(100.0)				
腎形成不全			1	13	44	197	130	17	1	403	16	419	3.81	0.40
(%)			(0.2)	(3.2)	(10.9)	(48.9)	(32.3)	(4.2)	(0.2)	(100.0)				
不明	14	104	403	1,318	4,866	8,922	3,198	214	12	19,051	1,074	20,125	3.56	0.47
(%)	(0.1)	(0.5)	(2.1)	(6.9)	(25.5)	(46.8)	(16.8)	(1.1)	(0.1)	(100.0)				
再導入	2	1	17	58	275	661	332	23	5	1,374	43	1,417	3.68	0.45
(%)	(0.1)	(0.1)	(1.2)	(4.2)	(20.0)	(48.1)	(24.2)	(1.7)	(0.4)	(100.0)				
その他	5	33	109	365	1,291	2,417	968	81	3	5,272	242	5,514	3.57	0.49
(%)	(0.1)	(0.6)	(2.1)	(6.9)	(24.5)	(45.8)	(18.4)	(1.5)	(0.1)	(100.0)				
合計	146	771	3,179	12,150	52,082	107,135	40,506	2,362	172	218,503	10,468	228,971	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(23.8)	(49.0)	(18.5)	(1.1)	(0.1)	(100.0)				
記載なし														
(%)														
総計	146	771	3,179	12,150	52,082	107,135	40,506	2,362	172	218,503	10,468	228,971	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(23.8)	(49.0)	(18.5)	(1.1)	(0.1)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査