

表 3247 透析前アルブミン濃度(g/dL) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	5	17	58	82	98	108	15	1	1	385	29	414	3.05	0.65
(%)	(1.3)	(4.4)	(15.1)	(21.3)	(25.5)	(28.1)	(3.9)	(0.3)	(0.3)	(100.0)				
私立大学	13	45	87	180	338	620	241	13	2	1,539	29	1,568	3.40	0.64
(%)	(0.8)	(2.9)	(5.7)	(11.7)	(22.0)	(40.3)	(15.7)	(0.8)	(0.1)	(100.0)				
国立	1	10	17	60	128	205	59	7	1	488	7	495	3.44	0.57
(%)	(0.2)	(2.0)	(3.5)	(12.3)	(26.2)	(42.0)	(12.1)	(1.4)	(0.2)	(100.0)				
県市町村立	19	101	327	932	3,266	5,422	1,700	105	13	11,885	945	12,830	3.50	0.49
(%)	(0.2)	(0.8)	(2.8)	(7.8)	(27.5)	(45.6)	(14.3)	(0.9)	(0.1)	(100.0)				
社会保険	5	26	81	219	814	1,200	377	18	2	2,742	54	2,796	3.48	0.49
(%)	(0.2)	(0.9)	(3.0)	(8.0)	(29.7)	(43.8)	(13.7)	(0.7)	(0.1)	(100.0)				
厚生連	8	40	122	459	1,791	2,795	1,045	73	4	6,337	251	6,588	3.54	0.48
(%)	(0.1)	(0.6)	(1.9)	(7.2)	(28.3)	(44.1)	(16.5)	(1.2)	(0.1)	(100.0)				
その他公的	7	38	157	524	1,862	3,248	1,044	63	4	6,947	318	7,265	3.53	0.47
(%)	(0.1)	(0.5)	(2.3)	(7.5)	(26.8)	(46.8)	(15.0)	(0.9)	(0.1)	(100.0)				
私立総合	10	26	106	340	1,256	2,282	806	37	2	4,865	119	4,984	3.54	0.47
(%)	(0.2)	(0.5)	(2.2)	(7.0)	(25.8)	(46.9)	(16.6)	(0.8)	(0.0)	(100.0)				
私立	66	362	1,604	5,359	17,096	28,256	9,663	567	58	63,031	3,568	66,599	3.51	0.49
(%)	(0.1)	(0.6)	(2.5)	(8.5)	(27.1)	(44.8)	(15.3)	(0.9)	(0.1)	(100.0)				
私立診療所	12	106	620	3,995	25,433	62,999	25,556	1,478	85	120,284	5,148	125,432	3.67	0.39
(%)	(0.0)	(0.1)	(0.5)	(3.3)	(21.1)	(52.4)	(21.2)	(1.2)	(0.1)	(100.0)				
総計	146	771	3,179	12,150	52,082	107,135	40,506	2,362	172	218,503	10,468	228,971	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(23.8)	(49.0)	(18.5)	(1.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査