

表 3251 透析前アルブミン濃度(g/dL) 透析前リン濃度(mg/dL)別 (施設血液透析患者 透析回数週3回)

透析前リン濃度	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
3未満	49	228	850	1,694	2,728	2,534	650	37	10	8,780	181	8,961	3.18	0.61
(%)	(0.6)	(2.6)	(9.7)	(19.3)	(31.1)	(28.9)	(7.4)	(0.4)	(0.1)	(100.0)				
3.0～	25	89	396	1,343	3,440	4,241	1,161	61	7	10,763	217	10,980	3.40	0.51
(%)	(0.2)	(0.8)	(3.7)	(12.5)	(32.0)	(39.4)	(10.8)	(0.6)	(0.1)	(100.0)				
3.5～	14	85	425	1,654	5,430	8,333	2,636	124	8	18,709	309	19,018	3.50	0.47
(%)	(0.1)	(0.5)	(2.3)	(8.8)	(29.0)	(44.5)	(14.1)	(0.7)	(0.0)	(100.0)				
4.0～	10	84	382	1,796	7,299	13,307	4,589	234	21	27,722	470	28,192	3.57	0.44
(%)	(0.0)	(0.3)	(1.4)	(6.5)	(26.3)	(48.0)	(16.6)	(0.8)	(0.1)	(100.0)				
4.5～	12	77	309	1,517	8,043	16,680	5,953	340	19	32,950	518	33,468	3.61	0.41
(%)	(0.0)	(0.2)	(0.9)	(4.6)	(24.4)	(50.6)	(18.1)	(1.0)	(0.1)	(100.0)				
5.0～	10	54	240	1,309	7,615	17,117	6,516	355	24	33,240	569	33,809	3.64	0.40
(%)	(0.0)	(0.2)	(0.7)	(3.9)	(22.9)	(51.5)	(19.6)	(1.1)	(0.1)	(100.0)				
5.5～	10	41	172	961	6,131	14,944	5,961	327	25	28,572	440	29,012	3.66	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.4)	(21.5)	(52.3)	(20.9)	(1.1)	(0.1)	(100.0)				
6.0～	1	7	25	162	1,042	2,528	1,005	75	6	4,851	77	4,928	3.67	0.39
(%)	(0.0)	(0.1)	(0.5)	(3.3)	(21.5)	(52.1)	(20.7)	(1.5)	(0.1)	(100.0)				
6.1～	3	19	75	516	3,313	8,544	3,555	203	13	16,241	264	16,505	3.68	0.39
(%)	(0.0)	(0.1)	(0.5)	(3.2)	(20.4)	(52.6)	(21.9)	(1.2)	(0.1)	(100.0)				
6.5～	5	22	77	423	2,756	7,399	3,216	196	15	14,109	268	14,377	3.69	0.39
(%)	(0.0)	(0.2)	(0.5)	(3.0)	(19.5)	(52.4)	(22.8)	(1.4)	(0.1)	(100.0)				
7.0～	1	9	52	259	1,622	4,593	1,983	161	6	8,686	174	8,860	3.70	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.0)	(18.7)	(52.9)	(22.8)	(1.9)	(0.1)	(100.0)				
7.5～	1	5	33	141	1,009	2,688	1,246	79	6	5,208	151	5,359	3.70	0.39
(%)	(0.0)	(0.1)	(0.6)	(2.7)	(19.4)	(51.6)	(23.9)	(1.5)	(0.1)	(100.0)				
8.0～		8	24	98	562	1,570	776	63	3	3,104	88	3,192	3.71	0.41
(%)		(0.3)	(0.8)	(3.2)	(18.1)	(50.6)	(25.0)	(2.0)	(0.1)	(100.0)				
8.5～		5	13	66	372	1,028	459	42	1	1,986	37	2,023	3.70	0.41
(%)		(0.3)	(0.7)	(3.3)	(18.7)	(51.8)	(23.1)	(2.1)	(0.1)	(100.0)				
9.0～	1	8	19	84	577	1,539	769	60	8	3,065	91	3,156	3.71	0.41
(%)	(0.0)	(0.3)	(0.6)	(2.7)	(18.8)	(50.2)	(25.1)	(2.0)	(0.3)	(100.0)				
合計	142	741	3,092	12,023	51,939	107,045	40,475	2,357	172	217,986	3,854	221,840	3.60	0.44
(%)	(0.1)	(0.3)	(1.4)	(5.5)	(23.8)	(49.1)	(18.6)	(1.1)	(0.1)	(100.0)				
記載なし	4	30	87	127	143	90	31	5		517	6,614	7,131	2.96	0.67
(%)	(0.8)	(5.8)	(16.8)	(24.6)	(27.7)	(17.4)	(6.0)	(1.0)		(100.0)				
総計	146	771	3,179	12,150	52,082	107,135	40,506	2,362	172	218,503	10,468	228,971	3.60	0.44
(%)	(0.1)	(0.4)	(1.5)	(5.6)	(23.8)	(49.0)	(18.5)	(1.1)	(0.1)	(100.0)				
平均	3.60	3.91	3.94	4.44	4.99	5.31	5.48	5.65	5.45	5.20	5.30	5.20		
標準偏差	1.71	1.75	1.63	1.49	1.40	1.36	1.39	1.47	1.73	1.42	1.58	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査