

表 3252 透析前アルブミン濃度(g/dL) PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:intact-PTH

PTH値	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
20未満	17	69	281	817	2,499	4,265	1,345	73	15	9,381	131	9,512	3.50	0.50
(%)	(0.2)	(0.7)	(3.0)	(8.7)	(26.6)	(45.5)	(14.3)	(0.8)	(0.2)	(100.0)				
20～	10	69	318	1,054	3,401	6,370	2,205	107	13	13,547	182	13,729	3.54	0.48
(%)	(0.1)	(0.5)	(2.3)	(7.8)	(25.1)	(47.0)	(16.3)	(0.8)	(0.1)	(100.0)				
40～	12	68	297	1,026	3,559	6,772	2,452	146	18	14,350	175	14,525	3.56	0.47
(%)	(0.1)	(0.5)	(2.1)	(7.1)	(24.8)	(47.2)	(17.1)	(1.0)	(0.1)	(100.0)				
60～	6	51	246	944	3,824	7,336	2,784	158	12	15,361	153	15,514	3.59	0.45
(%)	(0.0)	(0.3)	(1.6)	(6.1)	(24.9)	(47.8)	(18.1)	(1.0)	(0.1)	(100.0)				
80～	10	49	214	848	3,806	7,620	2,798	156	6	15,507	164	15,671	3.59	0.43
(%)	(0.1)	(0.3)	(1.4)	(5.5)	(24.5)	(49.1)	(18.0)	(1.0)	(0.0)	(100.0)				
100～	9	45	199	807	3,586	7,445	2,879	143	11	15,124	147	15,271	3.61	0.43
(%)	(0.1)	(0.3)	(1.3)	(5.3)	(23.7)	(49.2)	(19.0)	(0.9)	(0.1)	(100.0)				
120～	5	37	149	748	3,356	7,260	2,714	173	9	14,451	135	14,586	3.62	0.42
(%)	(0.0)	(0.3)	(1.0)	(5.2)	(23.2)	(50.2)	(18.8)	(1.2)	(0.1)	(100.0)				
140～	7	25	116	616	3,186	6,698	2,576	147	8	13,379	124	13,503	3.63	0.42
(%)	(0.1)	(0.2)	(0.9)	(4.6)	(23.8)	(50.1)	(19.3)	(1.1)	(0.1)	(100.0)				
160～	4	13	104	527	2,786	6,060	2,353	131	10	11,988	119	12,107	3.64	0.41
(%)	(0.0)	(0.1)	(0.9)	(4.4)	(23.2)	(50.6)	(19.6)	(1.1)	(0.1)	(100.0)				
180～	7	44	187	1,074	6,130	13,932	5,422	294	19	27,109	319	27,428	3.65	0.40
(%)	(0.0)	(0.2)	(0.7)	(4.0)	(22.6)	(51.4)	(20.0)	(1.1)	(0.1)	(100.0)				
240～			2	11	70	181	69	5		338		338	3.66	0.39
(%)			(0.6)	(3.3)	(20.7)	(53.6)	(20.4)	(1.5)		(100.0)				
241～	6	23	112	619	3,437	7,755	3,215	219	16	15,402	196	15,598	3.65	0.41
(%)	(0.0)	(0.1)	(0.7)	(4.0)	(22.3)	(50.4)	(20.9)	(1.4)	(0.1)	(100.0)				
300～	2	8	55	326	1,887	4,638	1,828	120	4	8,868	124	8,992	3.66	0.39
(%)	(0.0)	(0.1)	(0.6)	(3.7)	(21.3)	(52.3)	(20.6)	(1.4)	(0.0)	(100.0)				
360～	4	2	27	168	1,113	2,565	1,089	66	3	5,037	87	5,124	3.67	0.40
(%)	(0.1)	(0.0)	(0.5)	(3.3)	(22.1)	(50.9)	(21.6)	(1.3)	(0.1)	(100.0)				
420～		4	20	117	627	1,515	643	53	4	2,983	41	3,024	3.67	0.41
(%)		(0.1)	(0.7)	(3.9)	(21.0)	(50.8)	(21.6)	(1.8)	(0.1)	(100.0)				
480～	1	2	2	38	158	342	173	12	1	729	7	736	3.66	0.43
(%)	(0.1)	(0.3)	(0.3)	(5.2)	(21.7)	(46.9)	(23.7)	(1.6)	(0.1)	(100.0)				
500～	1	2	12	46	274	594	232	12	1	1,174	15	1,189	3.63	0.41
(%)	(0.1)	(0.2)	(1.0)	(3.9)	(23.3)	(50.6)	(19.8)	(1.0)	(0.1)	(100.0)				
540～			10	44	281	660	268	20	2	1,285	25	1,310	3.67	0.39
(%)			(0.8)	(3.4)	(21.9)	(51.4)	(20.9)	(1.6)	(0.2)	(100.0)				
600～		1	14	65	484	1,058	439	30	5	2,096	36	2,132	3.66	0.40
(%)		(0.0)	(0.7)	(3.1)	(23.1)	(50.5)	(20.9)	(1.4)	(0.2)	(100.0)				
800～		2	16	67	369	778	335	17		1,584	38	1,622	3.64	0.42
(%)		(0.1)	(1.0)	(4.2)	(23.3)	(49.1)	(21.1)	(1.1)		(100.0)				
合計	101	514	2,381	9,962	44,833	93,844	35,819	2,082	157	189,693	2,218	191,911	3.61	0.43
(%)	(0.1)	(0.3)	(1.3)	(5.3)	(23.6)	(49.5)	(18.9)	(1.1)	(0.1)	(100.0)				
記載なし	12	55	148	371	994	1,494	485	25	2	3,586	1,089	4,675	3.43	0.55
(%)	(0.3)	(1.5)	(4.1)	(10.3)	(27.7)	(41.7)	(13.5)	(0.7)	(0.1)	(100.0)				
総計	113	569	2,529	10,333	45,827	95,338	36,304	2,107	159	193,279	3,307	196,586	3.61	0.44
(%)	(0.1)	(0.3)	(1.3)	(5.3)	(23.7)	(49.3)	(18.8)	(1.1)	(0.1)	(100.0)				
平均	119.45	106.74	120.07	140.47	164.77	172.67	179.66	189.70	165.64	169.75	188.94	169.97		
標準偏差	109.85	111.70	141.18	147.03	166.89	158.13	159.67	187.36	154.19	160.37	205.17	160.97		

数値下のかつこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査