

表 3265 透析前アルブミン濃度(g/dL) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学		1	6	6	13	10	6			42		42	3.25	0.66
(%)		(2.4)	(14.3)	(14.3)	(31.0)	(23.8)	(14.3)			(100.0)				
私立大学		2	6	18	58	66	20			170	3	173	3.41	0.49
(%)		(1.2)	(3.5)	(10.6)	(34.1)	(38.8)	(11.8)			(100.0)				
国立		1	3	4	12	11	5			36		36	3.31	0.64
(%)		(2.8)	(8.3)	(11.1)	(33.3)	(30.6)	(13.9)			(100.0)				
県市町村立	1	8	28	100	480	838	224	10		1,689	84	1,773	3.53	0.43
(%)	(0.1)	(0.5)	(1.7)	(5.9)	(28.4)	(49.6)	(13.3)	(0.6)		(100.0)				
社会保険		1	2	18	88	153	47	1		310	9	319	3.57	0.41
(%)		(0.3)	(0.6)	(5.8)	(28.4)	(49.4)	(15.2)	(0.3)		(100.0)				
厚生連		3	15	61	264	457	133	10		943	70	1,013	3.55	0.43
(%)		(0.3)	(1.6)	(6.5)	(28.0)	(48.5)	(14.1)	(1.1)		(100.0)				
その他公的	2	1	10	40	170	333	91	9	1	657	37	694	3.57	0.44
(%)	(0.3)	(0.2)	(1.5)	(6.1)	(25.9)	(50.7)	(13.9)	(1.4)	(0.2)	(100.0)				
私立総合		4	15	73	301	526	152	8	1	1,080	13	1,093	3.55	0.44
(%)		(0.4)	(1.4)	(6.8)	(27.9)	(48.7)	(14.1)	(0.7)	(0.1)	(100.0)				
私立	3	25	139	520	2,513	4,878	1,487	42	7	9,614	401	10,015	3.57	0.41
(%)	(0.0)	(0.3)	(1.4)	(5.4)	(26.1)	(50.7)	(15.5)	(0.4)	(0.1)	(100.0)				
私立診療所	4	16	70	552	4,908	13,313	4,993	237	9	24,102	686	24,788	3.68	0.36
(%)	(0.0)	(0.1)	(0.3)	(2.3)	(20.4)	(55.2)	(20.7)	(1.0)	(0.0)	(100.0)				
総計	10	62	294	1,392	8,807	20,585	7,158	317	18	38,643	1,303	39,946	3.64	0.39
(%)	(0.0)	(0.2)	(0.8)	(3.6)	(22.8)	(53.3)	(18.5)	(0.8)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査