

表 3270 透析前アルブミン濃度(g/dL) PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

PTH値	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
20未満	1	10	28	94	447	817	269	5	1	1,672	11	1,683	3.55	0.44
(%)	(0.1)	(0.6)	(1.7)	(5.6)	(26.7)	(48.9)	(16.1)	(0.3)	(0.1)	(100.0)				
20～		6	35	114	491	1,162	341	20	1	2,170	25	2,195	3.60	0.42
(%)		(0.3)	(1.6)	(5.3)	(22.6)	(53.5)	(15.7)	(0.9)	(0.0)	(100.0)				
40～	3	3	30	99	580	1,266	431	13	1	2,426	11	2,437	3.61	0.40
(%)	(0.1)	(0.1)	(1.2)	(4.1)	(23.9)	(52.2)	(17.8)	(0.5)	(0.0)	(100.0)				
60～		7	26	111	613	1,299	497	27	1	2,581	11	2,592	3.63	0.41
(%)		(0.3)	(1.0)	(4.3)	(23.8)	(50.3)	(19.3)	(1.0)	(0.0)	(100.0)				
80～	1	6	20	124	702	1,458	490	25		2,826	20	2,846	3.61	0.40
(%)	(0.0)	(0.2)	(0.7)	(4.4)	(24.8)	(51.6)	(17.3)	(0.9)		(100.0)				
100～	1	2	24	114	656	1,420	529	23	3	2,772	23	2,795	3.63	0.40
(%)	(0.0)	(0.1)	(0.9)	(4.1)	(23.7)	(51.2)	(19.1)	(0.8)	(0.1)	(100.0)				
120～		3	16	86	579	1,345	516	20		2,565	16	2,581	3.65	0.38
(%)		(0.1)	(0.6)	(3.4)	(22.6)	(52.4)	(20.1)	(0.8)		(100.0)				
140～	1	1	11	64	492	1,276	444	21		2,310	13	2,323	3.66	0.36
(%)	(0.0)	(0.0)	(0.5)	(2.8)	(21.3)	(55.2)	(19.2)	(0.9)		(100.0)				
160～		3	7	65	464	1,179	404	16	1	2,139	11	2,150	3.65	0.37
(%)		(0.1)	(0.3)	(3.0)	(21.7)	(55.1)	(18.9)	(0.7)	(0.0)	(100.0)				
180～	1	7	11	130	1,046	2,729	1,002	49	1	4,976	29	5,005	3.67	0.36
(%)	(0.0)	(0.1)	(0.2)	(2.6)	(21.0)	(54.8)	(20.1)	(1.0)	(0.0)	(100.0)				
240～			1	3	18	41	14			77		77	3.59	0.41
(%)			(1.3)	(3.9)	(23.4)	(53.2)	(18.2)			(100.0)				
241～	1	2	11	61	612	1,641	574	22	3	2,927	28	2,955	3.68	0.36
(%)	(0.0)	(0.1)	(0.4)	(2.1)	(20.9)	(56.1)	(19.6)	(0.8)	(0.1)	(100.0)				
300～	1	2	8	47	343	943	334	15	3	1,696	18	1,714	3.67	0.38
(%)	(0.1)	(0.1)	(0.5)	(2.8)	(20.2)	(55.6)	(19.7)	(0.9)	(0.2)	(100.0)				
360～			3	27	228	551	210	11	2	1,032	10	1,042	3.67	0.37
(%)			(0.3)	(2.6)	(22.1)	(53.4)	(20.3)	(1.1)	(0.2)	(100.0)				
420～			3	5	121	323	130	6	1	589	2	591	3.71	0.34
(%)			(0.5)	(0.8)	(20.5)	(54.8)	(22.1)	(1.0)	(0.2)	(100.0)				
480～			1	7	29	80	26			143	3	146	3.63	0.38
(%)			(0.7)	(4.9)	(20.3)	(55.9)	(18.2)			(100.0)				
500～			7	44	127	57	57	2		237		237	3.70	0.36
(%)			(3.0)	(18.6)	(53.6)	(24.1)	(0.8)			(100.0)				
540～		1	5	46	143	49	49	3		247	5	252	3.69	0.35
(%)		(0.4)	(2.0)	(18.6)	(57.9)	(19.8)	(1.2)			(100.0)				
600～		3	7	83	245	75	75	7		420	3	423	3.68	0.36
(%)		(0.7)	(1.7)	(19.8)	(58.3)	(17.9)	(1.7)			(100.0)				
800～		5	13	64	198	66	66	1		347	3	350	3.64	0.37
(%)		(1.4)	(3.7)	(18.4)	(57.1)	(19.0)	(0.3)			(100.0)				
合計	10	52	244	1,183	7,658	18,243	6,458	286	18	34,152	242	34,394	3.64	0.39
(%)	(0.0)	(0.2)	(0.7)	(3.5)	(22.4)	(53.4)	(18.9)	(0.8)	(0.1)	(100.0)				
記載なし		4	11	67	180	229	74	2		567	167	734	3.45	0.47
(%)		(0.7)	(1.9)	(11.8)	(31.7)	(40.4)	(13.1)	(0.4)		(100.0)				
総計	10	56	255	1,250	7,838	18,472	6,532	288	18	34,719	409	35,128	3.64	0.39
(%)	(0.0)	(0.2)	(0.7)	(3.6)	(22.6)	(53.2)	(18.8)	(0.8)	(0.1)	(100.0)				
平均	132.30	103.23	133.94	147.07	168.69	181.16	183.14	184.95	211.22	177.13	194.48	177.26		
標準偏差	104.34	86.94	179.03	172.01	166.37	171.10	166.60	153.33	138.28	169.26	194.21	169.45		

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査