

表 3282 アルブミン濃度(g/dL) 原疾患別（腹膜透析患者）

原疾患	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	5	13	58	308	772	728	170	6		2,060	999	3,059	3.35	0.48
(%)	(0.2)	(0.6)	(2.8)	(15.0)	(37.5)	(35.3)	(8.3)	(0.3)		(100.0)				
慢性腎盂腎炎	1		4	10	13	27	11	1		67	26	93	3.41	0.61
(%)	(1.5)		(6.0)	(14.9)	(19.4)	(40.3)	(16.4)	(1.5)		(100.0)				
急速進行性糸球体腎炎		1	3	8	14	18	3	1		48	26	74	3.30	0.60
(%)		(2.1)	(6.3)	(16.7)	(29.2)	(37.5)	(6.3)	(2.1)		(100.0)				
妊娠腎/妊娠中毒症				2	8	8	5			23	9	32	3.56	0.50
(%)				(8.7)	(34.8)	(34.8)	(21.7)			(100.0)				
その他の分類不能の腎炎		1	3	7	14	19	13			57	28	85	3.47	0.60
(%)		(1.8)	(5.3)	(12.3)	(24.6)	(33.3)	(22.8)			(100.0)				
多発性嚢胞腎		7	10	34	56	36	13			156	87	243	3.16	0.55
(%)		(4.5)	(6.4)	(21.8)	(35.9)	(23.1)	(8.3)			(100.0)				
腎硬化症	1	4	46	139	270	240	37	2		739	344	1,083	3.24	0.49
(%)	(0.1)	(0.5)	(6.2)	(18.8)	(36.5)	(32.5)	(5.0)	(0.3)		(100.0)				
悪性高血圧	1			10	17	29	11			68	39	107	3.47	0.52
(%)	(1.5)			(14.7)	(25.0)	(42.6)	(16.2)			(100.0)				
糖尿病性腎症	3	30	118	369	662	510	109	4		1,805	961	2,766	3.20	0.53
(%)	(0.2)	(1.7)	(6.5)	(20.4)	(36.7)	(28.3)	(6.0)	(0.2)		(100.0)				
SLE腎炎		1	4	5	15	17				42	20	62	3.22	0.52
(%)		(2.4)	(9.5)	(11.9)	(35.7)	(40.5)				(100.0)				
アミロイド腎					4					4	5	9	3.30	0.12
(%)					(100.0)					(100.0)				
痛風腎				2	12	10	1			25	8	33	3.41	0.32
(%)				(8.0)	(48.0)	(40.0)	(4.0)			(100.0)				
先天性代謝異常による腎不全				3	8	3				14	4	18	3.26	0.37
(%)				(21.4)	(57.1)	(21.4)				(100.0)				
腎・尿路結核						1				1		1	3.70	
(%)						(100.0)				(100.0)				
腎・尿路結石			1	1	5	3				10	6	16	3.18	0.51
(%)			(10.0)	(10.0)	(50.0)	(30.0)				(100.0)				
腎・尿路腫瘍				4	3	1	1			9	5	14	3.10	0.52
(%)				(44.4)	(33.3)	(11.1)	(11.1)			(100.0)				
閉塞性尿路障害				3	5	5	2		1	16	6	22	3.54	0.72
(%)				(18.8)	(31.3)	(31.3)	(12.5)		(6.3)	(100.0)				
骨髄腫		1		2		2				5	3	8	3.10	0.72
(%)		(20.0)		(40.0)		(40.0)				(100.0)				
腎形成不全			1	2	22	15	10	1		51	25	76	3.52	0.48
(%)			(2.0)	(3.9)	(43.1)	(29.4)	(19.6)	(2.0)		(100.0)				
不明	4	8	35	90	193	179	26	5	1	541	299	840	3.24	0.56
(%)	(0.7)	(1.5)	(6.5)	(16.6)	(35.7)	(33.1)	(4.8)	(0.9)	(0.2)	(100.0)				
再導入			1	5	17	9	4			36	15	51	3.35	0.44
(%)			(2.8)	(13.9)	(47.2)	(25.0)	(11.1)			(100.0)				
その他		6	13	33	48	46	15	1		162	87	249	3.21	0.59
(%)		(3.7)	(8.0)	(20.4)	(29.6)	(28.4)	(9.3)	(0.6)		(100.0)				
合計	15	71	298	1,037	2,158	1,906	431	21	2	5,939	3,002	8,941	3.28	0.52
(%)	(0.3)	(1.2)	(5.0)	(17.5)	(36.3)	(32.1)	(7.3)	(0.4)	(0.0)	(100.0)				
記載なし														
(%)														
総計	15	71	298	1,037	2,158	1,906	431	21	2	5,939	3,002	8,941	3.28	0.52
(%)	(0.3)	(1.2)	(5.0)	(17.5)	(36.3)	(32.1)	(7.3)	(0.4)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査