

表 3283 アルブミン濃度(g/dL) 医療機関別（腹膜透析患者）

医療機関	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
国公立大学		3	30	62	181	185	36	4		501	212	713	3.34	0.50
(%)		(0.6)	(6.0)	(12.4)	(36.1)	(36.9)	(7.2)	(0.8)		(100.0)				
私立大学		5	18	113	237	227	62	7		669	464	1,133	3.35	0.50
(%)		(0.7)	(2.7)	(16.9)	(35.4)	(33.9)	(9.3)	(1.0)		(100.0)				
国立		4	8	37	51	39	9			148	54	202	3.16	0.54
(%)		(2.7)	(5.4)	(25.0)	(34.5)	(26.4)	(6.1)			(100.0)				
県市町村立	2	13	60	187	362	311	76	3		1,014	572	1,586	3.26	0.53
(%)	(0.2)	(1.3)	(5.9)	(18.4)	(35.7)	(30.7)	(7.5)	(0.3)		(100.0)				
社会保険		3	13	28	88	57	11	1		201	286	487	3.23	0.50
(%)		(1.5)	(6.5)	(13.9)	(43.8)	(28.4)	(5.5)	(0.5)		(100.0)				
厚生連	2	7	21	114	188	115	29			476	150	626	3.19	0.51
(%)	(0.4)	(1.5)	(4.4)	(23.9)	(39.5)	(24.2)	(6.1)			(100.0)				
その他公的	2	11	47	162	340	306	65	1	1	935	384	1,319	3.26	0.51
(%)	(0.2)	(1.2)	(5.0)	(17.3)	(36.4)	(32.7)	(7.0)	(0.1)	(0.1)	(100.0)				
私立総合	1	3	15	48	92	59	9			227	150	377	3.19	0.51
(%)	(0.4)	(1.3)	(6.6)	(21.1)	(40.5)	(26.0)	(4.0)			(100.0)				
私立	6	21	65	192	358	368	87	4	1	1,102	587	1,689	3.26	0.56
(%)	(0.5)	(1.9)	(5.9)	(17.4)	(32.5)	(33.4)	(7.9)	(0.4)	(0.1)	(100.0)				
私立診療所	2	1	21	94	261	239	47	1		666	143	809	3.35	0.46
(%)	(0.3)	(0.2)	(3.2)	(14.1)	(39.2)	(35.9)	(7.1)	(0.2)		(100.0)				
総計	15	71	298	1,037	2,158	1,906	431	21	2	5,939	3,002	8,941	3.28	0.52
(%)	(0.3)	(1.2)	(5.0)	(17.5)	(36.3)	(32.1)	(7.3)	(0.4)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査