

表 3284 アルブミン濃度(g/dL) 体重(kg)別 (腹膜透析患者)

体重	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
20未満		1	1	9	12	7	5	1		36	2	38	3.29	0.57
(%)		(2.8)	(2.8)	(25.0)	(33.3)	(19.4)	(13.9)	(2.8)		(100.0)				
20～		1	2	7	7	3	1			21	2	23	3.03	0.62
(%)		(4.8)	(9.5)	(33.3)	(33.3)	(14.3)	(4.8)			(100.0)				
30～	3	6	27	45	45	35	6			167	5	172	2.96	0.63
(%)	(1.8)	(3.6)	(16.2)	(26.9)	(26.9)	(21.0)	(3.6)			(100.0)				
40～	3	20	62	183	304	248	48	1	1	870	34	904	3.18	0.55
(%)	(0.3)	(2.3)	(7.1)	(21.0)	(34.9)	(28.5)	(5.5)	(0.1)	(0.1)	(100.0)				
50～	2	18	93	323	559	449	83	5	1	1,533	58	1,591	3.22	0.51
(%)	(0.1)	(1.2)	(6.1)	(21.1)	(36.5)	(29.3)	(5.4)	(0.3)	(0.1)	(100.0)				
60～	1	9	45	230	618	541	103	5		1,552	64	1,616	3.33	0.46
(%)	(0.1)	(0.6)	(2.9)	(14.8)	(39.8)	(34.9)	(6.6)	(0.3)		(100.0)				
70～	1	3	17	102	290	286	76	5		780	31	811	3.39	0.47
(%)	(0.1)	(0.4)	(2.2)	(13.1)	(37.2)	(36.7)	(9.7)	(0.6)		(100.0)				
80～			7	31	89	113	42	2		284	14	298	3.48	0.46
(%)			(2.5)	(10.9)	(31.3)	(39.8)	(14.8)	(0.7)		(100.0)				
90～	1	2	2	11	31	39	14			100	4	104	3.39	0.54
(%)	(1.0)	(2.0)	(2.0)	(11.0)	(31.0)	(39.0)	(14.0)			(100.0)				
100～				1	6	9	3			19	1	20	3.64	0.37
(%)				(5.3)	(31.6)	(47.4)	(15.8)			(100.0)				
110～					5	4				9	1	10	3.48	0.27
(%)					(55.6)	(44.4)				(100.0)				
120～			1		4	3	1			9		9	3.38	0.58
(%)			(11.1)		(44.4)	(33.3)	(11.1)			(100.0)				
合計	11	60	257	942	1,970	1,737	382	19	2	5,380	216	5,596	3.28	0.51
(%)	(0.2)	(1.1)	(4.8)	(17.5)	(36.6)	(32.3)	(7.1)	(0.4)	(0.0)	(100.0)				
記載なし	4	11	41	95	188	169	49	2		559	2,786	3,345	3.23	0.59
(%)	(0.7)	(2.0)	(7.3)	(17.0)	(33.6)	(30.2)	(8.8)	(0.4)		(100.0)				
総計	15	71	298	1,037	2,158	1,906	431	21	2	5,939	3,002	8,941	3.28	0.52
(%)	(0.3)	(1.2)	(5.0)	(17.5)	(36.3)	(32.1)	(7.3)	(0.4)	(0.0)	(100.0)				
平均	52.49	52.11	54.69	57.44	60.95	62.61	64.74	63.31	46.95	60.73	61.14	60.74		
標準偏差	19.30	13.47	12.92	12.93	13.43	13.67	16.08	17.14	6.86	13.88	14.41	13.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査