

表 3288 アルブミン濃度(g/dL) PTH値(pg/mL)別（腹膜透析患者）PTH測定法:intact-PTH

PTH値	1.5未満	1.5～	2.0～	2.5～	3.0～	3.5～	4.0～	4.5～	5.0～	合計	記載なし	総計	平均	標準偏差
20未満		2	10	25	41	32	8	2		120	3	123	3.24	0.56
(%)		(1.7)	(8.3)	(20.8)	(34.2)	(26.7)	(6.7)	(1.7)		(100.0)				
20～	3	2	20	44	113	78	19			279	2	281	3.22	0.53
(%)	(1.1)	(0.7)	(7.2)	(15.8)	(40.5)	(28.0)	(6.8)			(100.0)				
40～	1	2	10	45	92	92	19	2		263	5	268	3.31	0.52
(%)	(0.4)	(0.8)	(3.8)	(17.1)	(35.0)	(35.0)	(7.2)	(0.8)		(100.0)				
60～		5	16	44	113	110	18	2		308	7	315	3.29	0.51
(%)		(1.6)	(5.2)	(14.3)	(36.7)	(35.7)	(5.8)	(0.6)		(100.0)				
80～	1	3	8	53	117	102	21	1		306	3	309	3.31	0.48
(%)	(0.3)	(1.0)	(2.6)	(17.3)	(38.2)	(33.3)	(6.9)	(0.3)		(100.0)				
100～		4	10	62	115	103	12	2	1	309	7	316	3.27	0.49
(%)		(1.3)	(3.2)	(20.1)	(37.2)	(33.3)	(3.9)	(0.6)	(0.3)	(100.0)				
120～		2	18	49	116	91	23	2		301	4	305	3.28	0.51
(%)		(0.7)	(6.0)	(16.3)	(38.5)	(30.2)	(7.6)	(0.7)		(100.0)				
140～	1		10	45	103	107	18			284	1	285	3.32	0.47
(%)	(0.4)		(3.5)	(15.8)	(36.3)	(37.7)	(6.3)			(100.0)				
160～		2	16	48	92	101	14			273	3	276	3.28	0.50
(%)		(0.7)	(5.9)	(17.6)	(33.7)	(37.0)	(5.1)			(100.0)				
180～	1	11	32	105	260	178	49	3	1	640	14	654	3.27	0.53
(%)	(0.2)	(1.7)	(5.0)	(16.4)	(40.6)	(27.8)	(7.7)	(0.5)	(0.2)	(100.0)				
240～	1		1		6	1	1			10		10	3.07	0.83
(%)	(10.0)		(10.0)		(60.0)	(10.0)	(10.0)			(100.0)				
241～	1	2	22	66	157	142	37	1		428	3	431	3.32	0.49
(%)	(0.2)	(0.5)	(5.1)	(15.4)	(36.7)	(33.2)	(8.6)	(0.2)		(100.0)				
300～		5	13	58	110	110	34			330	1	331	3.31	0.53
(%)		(1.5)	(3.9)	(17.6)	(33.3)	(33.3)	(10.3)			(100.0)				
360～			10	33	71	59	22	1		196	4	200	3.33	0.49
(%)			(5.1)	(16.8)	(36.2)	(30.1)	(11.2)	(0.5)		(100.0)				
420～		1	7	29	54	36	12			139		139	3.25	0.51
(%)		(0.7)	(5.0)	(20.9)	(38.8)	(25.9)	(8.6)			(100.0)				
480～				8	12	20	2			42	1	43	3.39	0.40
(%)				(19.0)	(28.6)	(47.6)	(4.8)			(100.0)				
500～			3	12	15	21	3			54		54	3.27	0.51
(%)			(5.6)	(22.2)	(27.8)	(38.9)	(5.6)			(100.0)				
540～			4	9	25	31	3			72	1	73	3.35	0.50
(%)			(5.6)	(12.5)	(34.7)	(43.1)	(4.2)			(100.0)				
600～			5	19	43	29	7	1		104	1	105	3.28	0.46
(%)			(4.8)	(18.3)	(41.3)	(27.9)	(6.7)	(1.0)		(100.0)				
800～			2	20	46	31	8	1		108	2	110	3.29	0.46
(%)			(1.9)	(18.5)	(42.6)	(28.7)	(7.4)	(0.9)		(100.0)				
合計	9	41	217	774	1,701	1,474	330	18	2	4,566	62	4,628	3.29	0.51
(%)	(0.2)	(0.9)	(4.8)	(17.0)	(37.3)	(32.3)	(7.2)	(0.4)	(0.0)	(100.0)				
記載なし	3	11	17	40	60	47	15			193	183	376	3.08	0.65
(%)	(1.6)	(5.7)	(8.8)	(20.7)	(31.1)	(24.4)	(7.8)			(100.0)				
総計	12	52	234	814	1,761	1,521	345	18	2	4,759	245	5,004	3.28	0.51
(%)	(0.3)	(1.1)	(4.9)	(17.1)	(37.0)	(32.0)	(7.2)	(0.4)	(0.0)	(100.0)				
平均	119.56	163.59	208.01	230.59	225.78	220.67	239.36	206.61	173.50	224.21	198.92	223.88		
標準偏差	92.08	103.93	204.71	266.85	235.08	206.46	235.29	223.15	88.39	229.70	198.75	229.31		

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査