

表 3298 透析前CRP濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	110,049	36,786	18,887	12,260	7,575	2,720	1,351	739	881	595	191,843	63,798	255,641	0.65	1.96
(%)	(57.4)	(19.2)	(9.8)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
Off-line HDF	2,705	953	511	367	249	92	36	20	27	23	4,983	1,332	6,315	0.75	2.11
(%)	(54.3)	(19.1)	(10.3)	(7.4)	(5.0)	(1.8)	(0.7)	(0.4)	(0.5)	(0.5)	(100.0)				
On-line HDF	18,095	5,554	2,686	1,659	896	323	135	79	82	67	29,576	6,514	36,090	0.52	1.72
(%)	(61.2)	(18.8)	(9.1)	(5.6)	(3.0)	(1.1)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
Push/Pull HDF	235	56	29	22	17	3	2	1			365	172	537	0.46	1.01
(%)	(64.4)	(15.3)	(7.9)	(6.0)	(4.7)	(0.8)	(0.5)	(0.3)			(100.0)				
ハイオフィルトレーション	148	75	30	22	19	3	3	1	1		302	39	341	0.65	1.33
(%)	(49.0)	(24.8)	(9.9)	(7.3)	(6.3)	(1.0)	(1.0)	(0.3)	(0.3)		(100.0)				
血液濾過	45	15	4	6					1		71	21	92	0.46	1.69
(%)	(63.4)	(21.1)	(5.6)	(8.5)					(1.4)		(100.0)				
血液吸着透析 (リクセル)	755	298	132	95	39	17	8	6	8	1	1,359	264	1,623	0.59	1.58
(%)	(55.6)	(21.9)	(9.7)	(7.0)	(2.9)	(1.3)	(0.6)	(0.4)	(0.6)	(0.1)	(100.0)				
血液吸着透析 (その他)	2	1									3	4	7	0.20	0.24
(%)	(66.7)	(33.3)									(100.0)				
在宅血液透析	208	74	29	9	8	3	1				332	189	521	0.35	0.77
(%)	(62.7)	(22.3)	(8.7)	(2.7)	(2.4)	(0.9)	(0.3)				(100.0)				
腹膜透析 (CAPD)	1,636	533	265	163	112	48	19	13	22	4	2,815	2,206	5,021	0.65	1.77
(%)	(58.1)	(18.9)	(9.4)	(5.8)	(4.0)	(1.7)	(0.7)	(0.5)	(0.8)	(0.1)	(100.0)				
腹膜透析 (APD)	1,011	319	142	102	68	22	15	7	9	3	1,698	1,141	2,839	0.61	1.73
(%)	(59.5)	(18.8)	(8.4)	(6.0)	(4.0)	(1.3)	(0.9)	(0.4)	(0.5)	(0.2)	(100.0)				
腹膜透析 (CCPD)	420	143	42	43	22	13	6	4	3		696	385	1,081	0.57	1.40
(%)	(60.3)	(20.5)	(6.0)	(6.2)	(3.2)	(1.9)	(0.9)	(0.6)	(0.4)		(100.0)				
合計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし															
(%)															
総計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査