

表 3299 透析前CRP濃度(mg/dL) 併用の状況別 (透析患者全体)

| 併用の状況              | 0.2未満   | 0.2～    | 0.5～   | 1.0～   | 2.0～  | 4.0～  | 6.0～  | 8.0～  | 10.0～ | 15.0～ | 合計      | 記載なし   | 総計      | 平均   | 標準偏差 |
|--------------------|---------|---------|--------|--------|-------|-------|-------|-------|-------|-------|---------|--------|---------|------|------|
| HD(F)単独            | 132,088 | 43,753  | 22,277 | 14,426 | 8,788 | 3,154 | 1,535 | 846   | 998   | 681   | 228,546 | 72,248 | 300,794 | 0.63 | 1.92 |
| (%)                | (57.8)  | (19.1)  | (9.7)  | (6.3)  | (3.8) | (1.4) | (0.7) | (0.4) | (0.4) | (0.3) | (100.0) |        |         |      |      |
| PD単独               | 2,382   | 772     | 339    | 246    | 153   | 59    | 32    | 18    | 24    | 7     | 4,032   | 3,156  | 7,188   | 0.62 | 1.73 |
| (%)                | (59.1)  | (19.1)  | (8.4)  | (6.1)  | (3.8) | (1.5) | (0.8) | (0.4) | (0.6) | (0.2) | (100.0) |        |         |      |      |
| HD(F)単独.PD力テ<br>あり | 109     | 38      | 20     | 12     | 11    | 4     | 1     |       | 2     | 3     | 200     | 69     | 269     | 0.99 | 3.04 |
| (%)                | (54.5)  | (19.0)  | (10.0) | (6.0)  | (5.5) | (2.0) | (0.5) |       | (1.0) | (1.5) | (100.0) |        |         |      |      |
| PD.HD併用週1<br>HD(F) | 609     | 198     | 94     | 50     | 42    | 21    | 8     | 4     | 6     |       | 1,032   | 512    | 1,544   | 0.61 | 1.51 |
| (%)                | (59.0)  | (19.2)  | (9.1)  | (4.8)  | (4.1) | (2.0) | (0.8) | (0.4) | (0.6) |       | (100.0) |        |         |      |      |
| PD.HD併用週2<br>HD(F) | 65      | 25      | 16     | 11     | 5     | 3     |       | 1     | 2     |       | 128     | 49     | 177     | 0.84 | 2.02 |
| (%)                | (50.8)  | (19.5)  | (12.5) | (8.6)  | (3.9) | (2.3) |       | (0.8) | (1.6) |       | (100.0) |        |         |      |      |
| PD.HD併用週3<br>HD(F) | 12      | 11      | 3      | 1      | 3     | 3     |       |       |       | 1     | 34      | 6      | 40      | 1.44 | 3.42 |
| (%)                | (35.3)  | (32.4)  | (8.8)  | (2.9)  | (8.8) | (8.8) |       |       |       | (2.9) | (100.0) |        |         |      |      |
| PD.HD併用週4<br>HD(F) |         | 1       |        |        |       |       |       |       |       |       | 1       |        | 1       | 0.43 |      |
| (%)                |         | (100.0) |        |        |       |       |       |       |       |       | (100.0) |        |         |      |      |
| PD.HD併用その他         | 36      | 7       | 5      | 2      | 3     |       |       | 1     | 2     |       | 56      | 17     | 73      | 0.94 | 2.48 |
| (%)                | (64.3)  | (12.5)  | (8.9)  | (3.6)  | (5.4) |       |       | (1.8) | (3.6) |       | (100.0) |        |         |      |      |
| 合計                 | 135,301 | 44,805  | 22,754 | 14,748 | 9,005 | 3,244 | 1,576 | 870   | 1,034 | 692   | 234,029 | 76,057 | 310,086 | 0.63 | 1.92 |
| (%)                | (57.8)  | (19.1)  | (9.7)  | (6.3)  | (3.8) | (1.4) | (0.7) | (0.4) | (0.4) | (0.3) | (100.0) |        |         |      |      |
| 不明                 | 8       | 2       | 3      |        |       |       |       |       |       | 1     | 14      | 8      | 22      | 1.61 | 5.17 |
| (%)                | (57.1)  | (14.3)  | (21.4) |        |       |       |       |       |       | (7.1) | (100.0) |        |         |      |      |
| 記載なし<br>(%)        |         |         |        |        |       |       |       |       |       |       |         |        |         |      |      |
| 総計                 | 135,309 | 44,807  | 22,757 | 14,748 | 9,005 | 3,244 | 1,576 | 870   | 1,034 | 693   | 234,043 | 76,065 | 310,108 | 0.63 | 1.92 |
| (%)                | (57.8)  | (19.1)  | (9.7)  | (6.3)  | (3.8) | (1.4) | (0.7) | (0.4) | (0.4) | (0.3) | (100.0) |        |         |      |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査