

表 3300 透析前CRP濃度(mg/dL) 性別（透析患者全体）

性別	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
男性	82,472	30,477	15,559	10,050	6,059	2,181	1,059	557	699	464	149,577	48,564	198,141	0.66	1.97
(%)	(55.1)	(20.4)	(10.4)	(6.7)	(4.1)	(1.5)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
女性	52,837	14,330	7,198	4,698	2,946	1,063	517	313	335	229	84,466	27,501	111,967	0.57	1.83
(%)	(62.6)	(17.0)	(8.5)	(5.6)	(3.5)	(1.3)	(0.6)	(0.4)	(0.4)	(0.3)	(100.0)				
合計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし															
(%)															
総計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査