

表 3303 透析前CRP濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	44,046	13,483	6,431	4,064	2,423	845	404	205	283	195	72,379	24,591	96,970	0.56	1.81
(%)	(60.9)	(18.6)	(8.9)	(5.6)	(3.3)	(1.2)	(0.6)	(0.3)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	1,368	433	215	132	95	32	22	17	11	6	2,331	711	3,042	0.66	1.84
(%)	(58.7)	(18.6)	(9.2)	(5.7)	(4.1)	(1.4)	(0.9)	(0.7)	(0.5)	(0.3)	(100.0)				
急速進行性糸球体腎炎	1,056	356	213	129	100	40	12	10	9	7	1,932	562	2,494	0.73	1.88
(%)	(54.7)	(18.4)	(11.0)	(6.7)	(5.2)	(2.1)	(0.6)	(0.5)	(0.5)	(0.4)	(100.0)				
妊娠腎/妊娠中毒症	808	205	89	63	29	8	6	6	3	2	1,219	395	1,614	0.47	1.70
(%)	(66.3)	(16.8)	(7.3)	(5.2)	(2.4)	(0.7)	(0.5)	(0.5)	(0.2)	(0.2)	(100.0)				
その他の分類不能の腎炎	657	190	100	57	31	8	6	4	2	4	1,059	314	1,373	0.54	1.81
(%)	(62.0)	(17.9)	(9.4)	(5.4)	(2.9)	(0.8)	(0.6)	(0.4)	(0.2)	(0.4)	(100.0)				
多発性嚢胞腎	4,708	1,684	883	527	313	117	52	29	59	28	8,400	2,606	11,006	0.67	2.09
(%)	(56.0)	(20.0)	(10.5)	(6.3)	(3.7)	(1.4)	(0.6)	(0.3)	(0.7)	(0.3)	(100.0)				
腎硬化症	12,114	4,349	2,269	1,424	913	315	143	95	85	57	21,764	6,534	28,298	0.65	1.90
(%)	(55.7)	(20.0)	(10.4)	(6.5)	(4.2)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
悪性高血圧	1,163	396	196	98	45	23	9	8	6	1	1,945	628	2,573	0.48	1.27
(%)	(59.8)	(20.4)	(10.1)	(5.0)	(2.3)	(1.2)	(0.5)	(0.4)	(0.3)	(0.1)	(100.0)				
糖尿病性腎症	50,732	17,458	9,017	5,997	3,625	1,276	633	357	406	277	89,778	28,303	118,081	0.66	1.93
(%)	(56.5)	(19.4)	(10.0)	(6.7)	(4.0)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
SLE腎炎	971	316	169	126	70	34	16	9	5	3	1,719	517	2,236	0.70	2.78
(%)	(56.5)	(18.4)	(9.8)	(7.3)	(4.1)	(2.0)	(0.9)	(0.5)	(0.3)	(0.2)	(100.0)				
アミロイド腎	183	68	41	25	18	6	2	3	3	2	351	100	451	0.85	2.24
(%)	(52.1)	(19.4)	(11.7)	(7.1)	(5.1)	(1.7)	(0.6)	(0.9)	(0.9)	(0.6)	(100.0)				
痛風腎	471	158	85	57	31	14	3	3	3	1	826	287	1,113	0.62	2.26
(%)	(57.0)	(19.1)	(10.3)	(6.9)	(3.8)	(1.7)	(0.4)	(0.4)	(0.4)	(0.1)	(100.0)				
先天性代謝異常による腎不全	129	38	9	12	7	5					200	72	272	0.44	0.94
(%)	(64.5)	(19.0)	(4.5)	(6.0)	(3.5)	(2.5)					(100.0)				
腎・尿路結核	80	26	19	8	13	5	1	1	3		156	50	206	0.94	1.95
(%)	(51.3)	(16.7)	(12.2)	(5.1)	(8.3)	(3.2)	(0.6)	(0.6)	(1.9)		(100.0)				
腎・尿路結石	250	78	54	29	21	9	2	1	1		445	136	581	0.58	1.21
(%)	(56.2)	(17.5)	(12.1)	(6.5)	(4.7)	(2.0)	(0.4)	(0.2)	(0.2)		(100.0)				
腎・尿路腫瘍	336	125	72	59	38	14	6	6	2	7	665	221	886	0.97	2.84
(%)	(50.5)	(18.8)	(10.8)	(8.9)	(5.7)	(2.1)	(0.9)	(0.9)	(0.3)	(1.1)	(100.0)				
閉塞性尿路障害	291	109	62	28	37	14	5	1	5	2	554	169	723	0.81	1.84
(%)	(52.5)	(19.7)	(11.2)	(5.1)	(6.7)	(2.5)	(0.9)	(0.2)	(0.9)	(0.4)	(100.0)				
骨髄腫	110	39	19	18	5	2	3	3	1	1	201	75	276	0.93	3.47
(%)	(54.7)	(19.4)	(9.5)	(9.0)	(2.5)	(1.0)	(1.5)	(1.5)	(0.5)	(0.5)	(100.0)				
腎形成不全	339	84	34	16	15	6		2		1	497	152	649	0.43	1.61
(%)	(68.2)	(16.9)	(6.8)	(3.2)	(3.0)	(1.2)		(0.4)		(0.2)	(100.0)				
不明	11,421	3,828	2,063	1,361	882	339	182	82	112	82	20,352	7,192	27,544	0.72	2.16
(%)	(56.1)	(18.8)	(10.1)	(6.7)	(4.3)	(1.7)	(0.9)	(0.4)	(0.6)	(0.4)	(100.0)				
再導入	964	329	147	108	69	21	8	2	7	4	1,659	504	2,163	0.58	1.70
(%)	(58.1)	(19.8)	(8.9)	(6.5)	(4.2)	(1.3)	(0.5)	(0.1)	(0.4)	(0.2)	(100.0)				
その他	3,112	1,055	570	410	225	111	61	26	28	13	5,611	1,946	7,557	0.71	1.90
(%)	(55.5)	(18.8)	(10.2)	(7.3)	(4.0)	(2.0)	(1.1)	(0.5)	(0.5)	(0.2)	(100.0)				
合計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし															
(%)															
総計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査