

表 3308 透析前CRP濃度(mg/dL) PTH値(pg/mL)別 (透析患者全体) PTH測定法:intact-PTH

PTH値	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	5,169	1,833	992	812	526	211	102	53	58	45	9,801	2,072	11,873	0.83	2.29
(%)	(52.7)	(18.7)	(10.1)	(8.3)	(5.4)	(2.2)	(1.0)	(0.5)	(0.6)	(0.5)	(100.0)				
20～	7,954	2,631	1,416	957	625	256	131	69	78	48	14,165	2,781	16,946	0.72	1.98
(%)	(56.2)	(18.6)	(10.0)	(6.8)	(4.4)	(1.8)	(0.9)	(0.5)	(0.6)	(0.3)	(100.0)				
40～	8,575	2,839	1,482	1,064	672	264	109	66	75	52	15,198	2,818	18,016	0.70	1.98
(%)	(56.4)	(18.7)	(9.8)	(7.0)	(4.4)	(1.7)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
60～	9,311	3,151	1,598	1,012	654	226	119	61	76	28	16,236	2,994	19,230	0.62	1.65
(%)	(57.3)	(19.4)	(9.8)	(6.2)	(4.0)	(1.4)	(0.7)	(0.4)	(0.5)	(0.2)	(100.0)				
80～	9,669	3,153	1,607	1,054	648	217	100	61	70	60	16,639	2,974	19,613	0.64	2.15
(%)	(58.1)	(18.9)	(9.7)	(6.3)	(3.9)	(1.3)	(0.6)	(0.4)	(0.4)	(0.4)	(100.0)				
100～	9,566	3,155	1,524	1,042	589	179	105	66	71	41	16,338	2,820	19,158	0.59	1.66
(%)	(58.6)	(19.3)	(9.3)	(6.4)	(3.6)	(1.1)	(0.6)	(0.4)	(0.4)	(0.3)	(100.0)				
120～	9,132	2,852	1,478	934	542	210	81	50	53	35	15,367	2,855	18,222	0.57	1.71
(%)	(59.4)	(18.6)	(9.6)	(6.1)	(3.5)	(1.4)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
140～	8,559	2,692	1,383	835	492	167	78	37	45	26	14,314	2,493	16,807	0.54	1.57
(%)	(59.8)	(18.8)	(9.7)	(5.8)	(3.4)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
160～	7,845	2,509	1,218	721	389	142	58	32	34	23	12,971	2,202	15,173	0.50	1.46
(%)	(60.5)	(19.3)	(9.4)	(5.6)	(3.0)	(1.1)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
180～	17,874	5,569	2,781	1,622	923	281	119	65	87	54	29,375	5,138	34,513	0.50	1.53
(%)	(60.8)	(19.0)	(9.5)	(5.5)	(3.1)	(1.0)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
240～	223	76	36	30	12	3	2		2	1	385	59	444	0.57	1.71
(%)	(57.9)	(19.7)	(9.4)	(7.8)	(3.1)	(0.8)	(0.5)		(0.5)	(0.3)	(100.0)				
241～	10,210	3,347	1,649	880	542	191	76	31	39	20	16,985	2,854	19,839	0.49	1.42
(%)	(60.1)	(19.7)	(9.7)	(5.2)	(3.2)	(1.1)	(0.4)	(0.2)	(0.2)	(0.1)	(100.0)				
300～	5,912	1,921	937	582	314	83	59	25	23	13	9,869	1,657	11,526	0.50	1.33
(%)	(59.9)	(19.5)	(9.5)	(5.9)	(3.2)	(0.8)	(0.6)	(0.3)	(0.2)	(0.1)	(100.0)				
360～	3,298	1,186	534	335	158	54	24	17	21	11	5,638	1,046	6,684	0.53	1.54
(%)	(58.5)	(21.0)	(9.5)	(5.9)	(2.8)	(1.0)	(0.4)	(0.3)	(0.4)	(0.2)	(100.0)				
420～	1,937	661	348	195	112	32	20	10	5	5	3,325	600	3,925	0.52	1.50
(%)	(58.3)	(19.9)	(10.5)	(5.9)	(3.4)	(1.0)	(0.6)	(0.3)	(0.2)	(0.2)	(100.0)				
480～	453	174	82	49	28	12	4		1	2	805	159	964	0.53	1.34
(%)	(56.3)	(21.6)	(10.2)	(6.1)	(3.5)	(1.5)	(0.5)		(0.1)	(0.2)	(100.0)				
500～	770	253	130	101	51	16	4	3	2	3	1,333	231	1,564	0.53	1.32
(%)	(57.8)	(19.0)	(9.8)	(7.6)	(3.8)	(1.2)	(0.3)	(0.2)	(0.2)	(0.2)	(100.0)				
540～	826	292	142	80	46	17	4	3		3	1,413	301	1,714	0.50	1.53
(%)	(58.5)	(20.7)	(10.0)	(5.7)	(3.3)	(1.2)	(0.3)	(0.2)		(0.2)	(100.0)				
600～	1,346	442	255	133	88	27	15	5	1	3	2,315	484	2,799	0.52	1.30
(%)	(58.1)	(19.1)	(11.0)	(5.7)	(3.8)	(1.2)	(0.6)	(0.2)	(0.0)	(0.1)	(100.0)				
800～	948	372	197	113	69	24	16	7	5	3	1,754	408	2,162	0.61	1.47
(%)	(54.0)	(21.2)	(11.2)	(6.4)	(3.9)	(1.4)	(0.9)	(0.4)	(0.3)	(0.2)	(100.0)				
合計	119,577	39,108	19,789	12,551	7,480	2,612	1,226	661	746	476	204,226	36,946	241,172	0.58	1.71
(%)	(58.6)	(19.1)	(9.7)	(6.1)	(3.7)	(1.3)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
記載なし	1,682	593	333	270	247	105	57	32	29	24	3,372	3,012	6,384	1.08	2.51
(%)	(49.9)	(17.6)	(9.9)	(8.0)	(7.3)	(3.1)	(1.7)	(0.9)	(0.9)	(0.7)	(100.0)				
総計	121,259	39,701	20,122	12,821	7,727	2,717	1,283	693	775	500	207,598	39,958	247,556	0.59	1.73
(%)	(58.4)	(19.1)	(9.7)	(6.2)	(3.7)	(1.3)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
平均	173.36	175.29	174.10	165.29	159.76	152.21	158.06	145.59	133.24	137.86	172.12	172.46	172.17		
標準偏差	157.72	167.40	168.54	172.57	162.70	167.59	228.25	170.81	129.75	144.54	162.44	172.02	163.94		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2014年調査