

表 3309 透析前CRP濃度(mg/dL) PTH値(pg/mL)別(透析患者全体) PTH測定法:whole-PTH

PTH値	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	898	340	191	120	99	40	14	5	20	13	1,740	154	1,894	0.99	3.53
(%)	(51.6)	(19.5)	(11.0)	(6.9)	(5.7)	(2.3)	(0.8)	(0.3)	(1.1)	(0.7)	(100.0)				
20～	1,219	399	217	141	88	35	17	11	21	10	2,158	173	2,331	0.83	3.24
(%)	(56.5)	(18.5)	(10.1)	(6.5)	(4.1)	(1.6)	(0.8)	(0.5)	(1.0)	(0.5)	(100.0)				
35～	1,055	340	193	138	77	34	10	11	15	11	1,884	159	2,043	0.81	2.55
(%)	(56.0)	(18.0)	(10.2)	(7.3)	(4.1)	(1.8)	(0.5)	(0.6)	(0.8)	(0.6)	(100.0)				
50～	1,622	552	284	191	107	46	18	16	26	20	2,882	271	3,153	0.86	3.43
(%)	(56.3)	(19.2)	(9.9)	(6.6)	(3.7)	(1.6)	(0.6)	(0.6)	(0.9)	(0.7)	(100.0)				
75～	1,354	507	268	155	102	38	17	12	16	14	2,483	265	2,748	0.83	3.25
(%)	(54.5)	(20.4)	(10.8)	(6.2)	(4.1)	(1.5)	(0.7)	(0.5)	(0.6)	(0.6)	(100.0)				
100～	1,187	406	222	137	72	29	7	12	12	11	2,095	203	2,298	0.73	2.62
(%)	(56.7)	(19.4)	(10.6)	(6.5)	(3.4)	(1.4)	(0.3)	(0.6)	(0.6)	(0.5)	(100.0)				
125～	815	268	146	74	41	16	15	8	7	6	1,396	143	1,539	0.69	2.38
(%)	(58.4)	(19.2)	(10.5)	(5.3)	(2.9)	(1.1)	(1.1)	(0.6)	(0.5)	(0.4)	(100.0)				
150～	24	8	3	3	1	1					40	8	48	0.45	0.98
(%)	(60.0)	(20.0)	(7.5)	(7.5)	(2.5)	(2.5)					(100.0)				
151～	629	263	85	73	31	10	6	2	9	4	1,112	151	1,263	0.59	1.73
(%)	(56.6)	(23.7)	(7.6)	(6.6)	(2.8)	(0.9)	(0.5)	(0.2)	(0.8)	(0.4)	(100.0)				
180～	443	161	76	33	28	9	4	5	6	6	771	104	875	0.78	2.71
(%)	(57.5)	(20.9)	(9.9)	(4.3)	(3.6)	(1.2)	(0.5)	(0.6)	(0.8)	(0.8)	(100.0)				
210～	288	130	43	48	21	8	2	3	5	4	552	76	628	0.95	4.51
(%)	(52.2)	(23.6)	(7.8)	(8.7)	(3.8)	(1.4)	(0.4)	(0.5)	(0.9)	(0.7)	(100.0)				
240～	358	111	52	36	19	4	3		3	3	589	85	674	0.59	1.97
(%)	(60.8)	(18.8)	(8.8)	(6.1)	(3.2)	(0.7)	(0.5)		(0.5)	(0.5)	(100.0)				
300～	143	55	22	18	8	2	1			3	252	32	284	0.69	2.56
(%)	(56.7)	(21.8)	(8.7)	(7.1)	(3.2)	(0.8)	(0.4)			(1.2)	(100.0)				
350～	94	26	23	6	7		1				157	22	179	0.42	0.84
(%)	(59.9)	(16.6)	(14.6)	(3.8)	(4.5)		(0.6)				(100.0)				
400～	93	37	21	12	7	2	1	2	2	3	180	22	202	1.20	3.96
(%)	(51.7)	(20.6)	(11.7)	(6.7)	(3.9)	(1.1)	(0.6)	(1.1)	(1.1)	(1.7)	(100.0)				
500～	122	42	16	15	12	2	1	1	1	4	216	31	247	1.05	3.66
(%)	(56.5)	(19.4)	(7.4)	(6.9)	(5.6)	(0.9)	(0.5)	(0.5)	(0.5)	(1.9)	(100.0)				
合計	10,344	3,645	1,862	1,200	720	276	117	88	143	112	18,507	1,899	20,406	0.80	3.02
(%)	(55.9)	(19.7)	(10.1)	(6.5)	(3.9)	(1.5)	(0.6)	(0.5)	(0.8)	(0.6)	(100.0)				
記載なし	80	28	14	10	6	5	2	1	2	2	150	507	657	1.20	3.30
(%)	(53.3)	(18.7)	(9.3)	(6.7)	(4.0)	(3.3)	(1.3)	(0.7)	(1.3)	(1.3)	(100.0)				
総計	10,424	3,673	1,876	1,210	726	281	119	89	145	114	18,657	2,406	21,063	0.81	3.03
(%)	(55.9)	(19.7)	(10.1)	(6.5)	(3.9)	(1.5)	(0.6)	(0.5)	(0.8)	(0.6)	(100.0)				
平均	107.53	108.41	102.43	105.78	104.54	87.43	95.02	98.64	88.99	134.41	106.56	122.11	108.01		
標準偏差	112.45	106.01	103.93	126.93	137.01	104.37	87.91	90.30	85.91	238.37	113.15	121.28	114.01		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査