

表 3314 透析前CRP濃度(mg/dL) 透析前収縮期血圧(mmHg)別 (透析患者全体)

透析前収縮期血圧	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
60未満	17	6	2	4	4				1	2	36	4	40	2.15	5.24
(%)	(47.2)	(16.7)	(5.6)	(11.1)	(11.1)				(2.8)	(5.6)	(100.0)				
60～	119	82	44	46	33	16	5	5	7	7	364	69	433	1.66	3.41
(%)	(32.7)	(22.5)	(12.1)	(12.6)	(9.1)	(4.4)	(1.4)	(1.4)	(1.9)	(1.9)	(100.0)				
80～	1,174	544	378	269	211	84	51	28	36	26	2,801	537	3,338	1.28	3.13
(%)	(41.9)	(19.4)	(13.5)	(9.6)	(7.5)	(3.0)	(1.8)	(1.0)	(1.3)	(0.9)	(100.0)				
100～	7,417	2,862	1,672	1,203	845	342	200	111	118	98	14,868	2,739	17,607	0.97	2.60
(%)	(49.9)	(19.2)	(11.2)	(8.1)	(5.7)	(2.3)	(1.3)	(0.7)	(0.8)	(0.7)	(100.0)				
120～	25,403	8,633	4,402	2,968	1,877	670	338	193	231	149	44,864	8,407	53,271	0.68	2.06
(%)	(56.6)	(19.2)	(9.8)	(6.6)	(4.2)	(1.5)	(0.8)	(0.4)	(0.5)	(0.3)	(100.0)				
140～	40,600	12,985	6,258	4,067	2,363	820	398	218	270	175	68,154	12,668	80,822	0.58	1.88
(%)	(59.6)	(19.1)	(9.2)	(6.0)	(3.5)	(1.2)	(0.6)	(0.3)	(0.4)	(0.3)	(100.0)				
160～	29,516	9,434	4,639	2,858	1,617	566	242	142	166	98	49,278	8,917	58,195	0.54	1.68
(%)	(59.9)	(19.1)	(9.4)	(5.8)	(3.3)	(1.1)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
180～	11,008	3,701	1,853	1,151	654	233	111	54	41	34	18,840	3,384	22,224	0.55	1.57
(%)	(58.4)	(19.6)	(9.8)	(6.1)	(3.5)	(1.2)	(0.6)	(0.3)	(0.2)	(0.2)	(100.0)				
200～	2,821	922	488	271	153	54	24	6	13	7	4,759	787	5,546	0.51	1.34
(%)	(59.3)	(19.4)	(10.3)	(5.7)	(3.2)	(1.1)	(0.5)	(0.1)	(0.3)	(0.1)	(100.0)				
220～	582	182	97	67	36	14	5			3	986	174	1,160	0.52	1.37
(%)	(59.0)	(18.5)	(9.8)	(6.8)	(3.7)	(1.4)	(0.5)			(0.3)	(100.0)				
合計	118,657	39,351	19,833	12,904	7,793	2,799	1,374	757	883	599	204,950	37,686	242,636	0.63	1.93
(%)	(57.9)	(19.2)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし	16,652	5,456	2,924	1,844	1,212	445	202	113	151	94	29,093	38,379	67,472	0.66	1.85
(%)	(57.2)	(18.8)	(10.1)	(6.3)	(4.2)	(1.5)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
総計	135,309	44,807	22,757	14,748	9,005	3,244	1,576	870	1,034	693	234,043	76,065	310,108	0.63	1.92
(%)	(57.8)	(19.1)	(9.7)	(6.3)	(3.8)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
平均	152.26	151.22	150.38	149.06	147.07	146.13	144.20	142.86	141.73	140.44	151.23	150.59	151.13		
標準偏差	23.54	24.33	25.23	25.56	26.10	26.74	26.71	25.69	25.47	27.54	24.26	24.05	24.23		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査