

表 3320 透析前CRP濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	32,757	10,216	4,912	3,075	1,870	627	320	161	216	150	54,304	14,273	68,577	0.57	1.81
(%)	(60.3)	(18.8)	(9.0)	(5.7)	(3.4)	(1.2)	(0.6)	(0.3)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	1,007	331	167	98	78	22	17	14	9	6	1,749	416	2,165	0.70	1.98
(%)	(57.6)	(18.9)	(9.5)	(5.6)	(4.5)	(1.3)	(1.0)	(0.8)	(0.5)	(0.3)	(100.0)				
急速進行性糸球体腎炎	810	291	168	108	87	33	9	8	9	6	1,529	305	1,834	0.78	1.99
(%)	(53.0)	(19.0)	(11.0)	(7.1)	(5.7)	(2.2)	(0.6)	(0.5)	(0.6)	(0.4)	(100.0)				
妊娠腎/妊娠中毒症	555	132	60	43	18	3	4	4	2	1	822	242	1,064	0.45	1.78
(%)	(67.5)	(16.1)	(7.3)	(5.2)	(2.2)	(0.4)	(0.5)	(0.5)	(0.2)	(0.1)	(100.0)				
その他の分類不能の腎炎	481	127	80	44	19	6	6	4	2	4	773	176	949	0.60	2.06
(%)	(62.2)	(16.4)	(10.3)	(5.7)	(2.5)	(0.8)	(0.8)	(0.5)	(0.3)	(0.5)	(100.0)				
多発性嚢胞腎	3,697	1,318	723	416	242	96	46	25	49	19	6,631	1,567	8,198	0.68	2.11
(%)	(55.8)	(19.9)	(10.9)	(6.3)	(3.6)	(1.4)	(0.7)	(0.4)	(0.7)	(0.3)	(100.0)				
腎硬化症	9,745	3,526	1,856	1,164	748	258	117	76	68	50	17,608	3,852	21,460	0.66	1.97
(%)	(55.3)	(20.0)	(10.5)	(6.6)	(4.2)	(1.5)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
悪性高血圧	899	305	150	77	37	20	7	7	5		1,507	374	1,881	0.49	1.27
(%)	(59.7)	(20.2)	(10.0)	(5.1)	(2.5)	(1.3)	(0.5)	(0.5)	(0.3)		(100.0)				
糖尿病性腎症	41,327	14,256	7,410	4,924	2,998	1,054	520	284	341	222	73,336	17,171	90,507	0.66	1.92
(%)	(56.4)	(19.4)	(10.1)	(6.7)	(4.1)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
SLE腎炎	706	242	125	93	55	27	11	7	5	3	1,274	276	1,550	0.75	3.16
(%)	(55.4)	(19.0)	(9.8)	(7.3)	(4.3)	(2.1)	(0.9)	(0.5)	(0.4)	(0.2)	(100.0)				
アミロイド腎	137	54	33	19	13	6	2	3	3	2	272	53	325	0.95	2.49
(%)	(50.4)	(19.9)	(12.1)	(7.0)	(4.8)	(2.2)	(0.7)	(1.1)	(1.1)	(0.7)	(100.0)				
痛風腎	366	122	68	44	23	11	2	3	3	1	643	185	828	0.66	2.51
(%)	(56.9)	(19.0)	(10.6)	(6.8)	(3.6)	(1.7)	(0.3)	(0.5)	(0.5)	(0.2)	(100.0)				
先天性代謝異常による腎不全	90	26	6	11	4	3					140	37	177	0.44	0.95
(%)	(64.3)	(18.6)	(4.3)	(7.9)	(2.9)	(2.1)					(100.0)				
腎・尿路結核	64	24	13	7	13	3	1	1	3		129	29	158	1.01	2.07
(%)	(49.6)	(18.6)	(10.1)	(5.4)	(10.1)	(2.3)	(0.8)	(0.8)	(2.3)		(100.0)				
腎・尿路結石	192	60	47	23	17	8	2	1	1		351	79	430	0.62	1.29
(%)	(54.7)	(17.1)	(13.4)	(6.6)	(4.8)	(2.3)	(0.6)	(0.3)	(0.3)		(100.0)				
腎・尿路腫瘍	284	104	56	51	30	14	6	5	2	5	557	129	686	0.93	2.44
(%)	(51.0)	(18.7)	(10.1)	(9.2)	(5.4)	(2.5)	(1.1)	(0.9)	(0.4)	(0.9)	(100.0)				
閉塞性尿路障害	219	90	51	24	30	9	5	1	3	2	434	83	517	0.82	1.86
(%)	(50.5)	(20.7)	(11.8)	(5.5)	(6.9)	(2.1)	(1.2)	(0.2)	(0.7)	(0.5)	(100.0)				
骨髄腫	80	31	13	14	4	1	2	2	1	1	149	41	190	1.00	3.91
(%)	(53.7)	(20.8)	(8.7)	(9.4)	(2.7)	(0.7)	(1.3)	(1.3)	(0.7)	(0.7)	(100.0)				
腎形成不全	234	57	24	10	9	6		1		1	342	77	419	0.46	1.84
(%)	(68.4)	(16.7)	(7.0)	(2.9)	(2.6)	(1.8)		(0.3)		(0.3)	(100.0)				
不明	8,777	3,010	1,629	1,121	733	270	150	64	91	72	15,917	4,208	20,125	0.75	2.20
(%)	(55.1)	(18.9)	(10.2)	(7.0)	(4.6)	(1.7)	(0.9)	(0.4)	(0.6)	(0.5)	(100.0)				
再導入	670	216	106	82	50	15	6	2	7	4	1,158	259	1,417	0.65	1.95
(%)	(57.9)	(18.7)	(9.2)	(7.1)	(4.3)	(1.3)	(0.5)	(0.2)	(0.6)	(0.3)	(100.0)				
その他	2,432	840	450	324	176	90	49	21	22	11	4,415	1,099	5,514	0.72	1.98
(%)	(55.1)	(19.0)	(10.2)	(7.3)	(4.0)	(2.0)	(1.1)	(0.5)	(0.5)	(0.2)	(100.0)				
合計	105,529	35,378	18,147	11,772	7,254	2,582	1,282	694	842	560	184,040	44,931	228,971	0.64	1.94
(%)	(57.3)	(19.2)	(9.9)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
記載なし															
(%)															
総計	105,529	35,378	18,147	11,772	7,254	2,582	1,282	694	842	560	184,040	44,931	228,971	0.64	1.94
(%)	(57.3)	(19.2)	(9.9)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査