

表 3323 透析前CRP濃度(mg/dL) 透析前補正カルシウム濃度(mg/dL)別(施設血液透析患者 透析回数週3回)

透析前補正 カルシウム濃度	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
7.0未満	416	136	60	46	27	7	6	10	1	1	710	164	874	0.63	1.64
(%)	(58.6)	(19.2)	(8.5)	(6.5)	(3.8)	(1.0)	(0.8)	(1.4)	(0.1)	(0.1)	(100.0)				
7.0～	595	207	115	54	25	8	4	3			1,011	255	1,266	0.42	0.92
(%)	(58.9)	(20.5)	(11.4)	(5.3)	(2.5)	(0.8)	(0.4)	(0.3)			(100.0)				
7.5～	2,421	838	425	212	122	36	16	5	17	9	4,101	870	4,971	0.52	2.00
(%)	(59.0)	(20.4)	(10.4)	(5.2)	(3.0)	(0.9)	(0.4)	(0.1)	(0.4)	(0.2)	(100.0)				
8.0～	8,220	2,762	1,259	764	356	100	55	21	37	21	13,595	2,695	16,290	0.47	1.55
(%)	(60.5)	(20.3)	(9.3)	(5.6)	(2.6)	(0.7)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
8.4～	2,404	737	359	212	110	31	12	8	12	7	3,892	784	4,676	0.47	1.34
(%)	(61.8)	(18.9)	(9.2)	(5.4)	(2.8)	(0.8)	(0.3)	(0.2)	(0.3)	(0.2)	(100.0)				
8.5～	27,428	8,672	4,214	2,534	1,386	459	170	110	123	81	45,177	8,725	53,902	0.50	1.58
(%)	(60.7)	(19.2)	(9.3)	(5.6)	(3.1)	(1.0)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
9.0～	31,977	10,582	5,353	3,374	1,989	684	332	161	177	125	54,754	10,482	65,236	0.58	1.75
(%)	(58.4)	(19.3)	(9.8)	(6.2)	(3.6)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
9.5～	19,701	6,788	3,523	2,512	1,606	564	298	165	182	98	35,437	6,967	42,404	0.70	1.94
(%)	(55.6)	(19.2)	(9.9)	(7.1)	(4.5)	(1.6)	(0.8)	(0.5)	(0.5)	(0.3)	(100.0)				
10.0～	2,301	801	483	301	230	90	35	26	39	13	4,319	908	5,227	0.81	2.12
(%)	(53.3)	(18.5)	(11.2)	(7.0)	(5.3)	(2.1)	(0.8)	(0.6)	(0.9)	(0.3)	(100.0)				
10.1～	5,593	2,051	1,191	841	639	240	135	80	102	74	10,946	2,331	13,277	0.95	2.42
(%)	(51.1)	(18.7)	(10.9)	(7.7)	(5.8)	(2.2)	(1.2)	(0.7)	(0.9)	(0.7)	(100.0)				
10.5～	2,479	955	571	462	373	172	79	43	63	57	5,254	1,199	6,453	1.23	3.32
(%)	(47.2)	(18.2)	(10.9)	(8.8)	(7.1)	(3.3)	(1.5)	(0.8)	(1.2)	(1.1)	(100.0)				
11.0～	708	306	199	167	173	70	48	20	30	26	1,747	418	2,165	1.60	3.40
(%)	(40.5)	(17.5)	(11.4)	(9.6)	(9.9)	(4.0)	(2.7)	(1.1)	(1.7)	(1.5)	(100.0)				
11.5～	199	114	90	86	57	26	14	11	12	12	621	148	769	1.80	3.66
(%)	(32.0)	(18.4)	(14.5)	(13.8)	(9.2)	(4.2)	(2.3)	(1.8)	(1.9)	(1.9)	(100.0)				
12.0～	132	73	66	52	50	26	25	5	12	11	452	87	539	2.24	4.00
(%)	(29.2)	(16.2)	(14.6)	(11.5)	(11.1)	(5.8)	(5.5)	(1.1)	(2.7)	(2.4)	(100.0)				
合計	104,574	35,022	17,908	11,617	7,143	2,513	1,229	668	807	535	182,016	36,033	218,049	0.63	1.90
(%)	(57.5)	(19.2)	(9.8)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし	955	356	239	155	111	69	53	26	35	25	2,024	8,898	10,922	1.44	4.18
(%)	(47.2)	(17.6)	(11.8)	(7.7)	(5.5)	(3.4)	(2.6)	(1.3)	(1.7)	(1.2)	(100.0)				
総計	105,529	35,378	18,147	11,772	7,254	2,582	1,282	694	842	560	184,040	44,931	228,971	0.64	1.94
(%)	(57.3)	(19.2)	(9.9)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
平均	9.13	9.17	9.22	9.29	9.40	9.48	9.56	9.51	9.60	9.67	9.18	9.20	9.18		
標準偏差	0.70	0.73	0.76	0.78	0.82	0.84	0.92	0.94	0.88	0.96	0.74	0.76	0.74		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査