

表 3325 透析前CRP濃度(mg/dL) PTH値(pg/mL)別(施設血液透析患者 透析回数週3回) PTH測定法:intact-PTH

PTH値	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	4,040	1,417	794	642	444	174	78	41	47	38	7,715	1,797	9,512	0.85	2.39
(%)	(52.4)	(18.4)	(10.3)	(8.3)	(5.8)	(2.3)	(1.0)	(0.5)	(0.6)	(0.5)	(100.0)				
20～	6,300	2,130	1,161	798	502	209	108	58	65	38	11,369	2,360	13,729	0.73	1.96
(%)	(55.4)	(18.7)	(10.2)	(7.0)	(4.4)	(1.8)	(0.9)	(0.5)	(0.6)	(0.3)	(100.0)				
40～	6,774	2,294	1,185	862	544	212	89	54	62	43	12,119	2,406	14,525	0.71	2.02
(%)	(55.9)	(18.9)	(9.8)	(7.1)	(4.5)	(1.7)	(0.7)	(0.4)	(0.5)	(0.4)	(100.0)				
60～	7,386	2,525	1,317	821	536	187	97	48	68	19	13,004	2,510	15,514	0.63	1.67
(%)	(56.8)	(19.4)	(10.1)	(6.3)	(4.1)	(1.4)	(0.7)	(0.4)	(0.5)	(0.1)	(100.0)				
80～	7,615	2,475	1,276	836	530	181	81	49	63	50	13,156	2,515	15,671	0.66	2.18
(%)	(57.9)	(18.8)	(9.7)	(6.4)	(4.0)	(1.4)	(0.6)	(0.4)	(0.5)	(0.4)	(100.0)				
100～	7,514	2,513	1,189	849	482	136	87	50	55	33	12,908	2,363	15,271	0.60	1.67
(%)	(58.2)	(19.5)	(9.2)	(6.6)	(3.7)	(1.1)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
120～	7,199	2,255	1,216	758	440	160	69	39	45	33	12,214	2,372	14,586	0.59	1.80
(%)	(58.9)	(18.5)	(10.0)	(6.2)	(3.6)	(1.3)	(0.6)	(0.3)	(0.4)	(0.3)	(100.0)				
140～	6,799	2,159	1,111	678	404	132	65	29	31	23	11,431	2,072	13,503	0.54	1.61
(%)	(59.5)	(18.9)	(9.7)	(5.9)	(3.5)	(1.2)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
160～	6,142	2,025	979	561	325	112	55	23	27	18	10,267	1,840	12,107	0.51	1.42
(%)	(59.8)	(19.7)	(9.5)	(5.5)	(3.2)	(1.1)	(0.5)	(0.2)	(0.3)	(0.2)	(100.0)				
180～	14,000	4,415	2,245	1,272	740	215	100	51	68	44	23,150	4,278	27,428	0.51	1.54
(%)	(60.5)	(19.1)	(9.7)	(5.5)	(3.2)	(0.9)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
240～	171	60	30	20	4	1	2		2	1	291	47	338	0.55	1.88
(%)	(58.8)	(20.6)	(10.3)	(6.9)	(1.4)	(0.3)	(0.7)		(0.7)	(0.3)	(100.0)				
241～	7,893	2,644	1,284	712	414	157	57	28	29	18	13,236	2,362	15,598	0.50	1.48
(%)	(59.6)	(20.0)	(9.7)	(5.4)	(3.1)	(1.2)	(0.4)	(0.2)	(0.2)	(0.1)	(100.0)				
300～	4,544	1,512	750	447	248	62	45	21	18	5	7,652	1,340	8,992	0.49	1.23
(%)	(59.4)	(19.8)	(9.8)	(5.8)	(3.2)	(0.8)	(0.6)	(0.3)	(0.2)	(0.1)	(100.0)				
360～	2,502	902	394	265	128	43	18	12	17	8	4,289	835	5,124	0.53	1.48
(%)	(58.3)	(21.0)	(9.2)	(6.2)	(3.0)	(1.0)	(0.4)	(0.3)	(0.4)	(0.2)	(100.0)				
420～	1,453	518	274	150	87	22	12	7	4	3	2,530	494	3,024	0.51	1.44
(%)	(57.4)	(20.5)	(10.8)	(5.9)	(3.4)	(0.9)	(0.5)	(0.3)	(0.2)	(0.1)	(100.0)				
480～	343	128	58	39	22	10	2		1	2	605	131	736	0.55	1.45
(%)	(56.7)	(21.2)	(9.6)	(6.4)	(3.6)	(1.7)	(0.3)		(0.2)	(0.3)	(100.0)				
500～	587	183	94	74	40	12	4	2	1	2	999	190	1,189	0.53	1.27
(%)	(58.8)	(18.3)	(9.4)	(7.4)	(4.0)	(1.2)	(0.4)	(0.2)	(0.1)	(0.2)	(100.0)				
540～	624	220	112	65	35	15	3	2		3	1,079	231	1,310	0.53	1.68
(%)	(57.8)	(20.4)	(10.4)	(6.0)	(3.2)	(1.4)	(0.3)	(0.2)		(0.3)	(100.0)				
600～	1,018	335	191	100	65	21	11	3	1	3	1,748	384	2,132	0.52	1.37
(%)	(58.2)	(19.2)	(10.9)	(5.7)	(3.7)	(1.2)	(0.6)	(0.2)	(0.1)	(0.2)	(100.0)				
800～	693	265	156	85	51	20	12	7	3	3	1,295	327	1,622	0.64	1.57
(%)	(53.5)	(20.5)	(12.0)	(6.6)	(3.9)	(1.5)	(0.9)	(0.5)	(0.2)	(0.2)	(100.0)				
合計	93,597	30,975	15,816	10,034	6,041	2,081	995	524	607	387	161,057	30,854	191,911	0.59	1.74
(%)	(58.1)	(19.2)	(9.8)	(6.2)	(3.8)	(1.3)	(0.6)	(0.3)	(0.4)	(0.2)	(100.0)				
記載なし	1,287	459	257	216	200	82	44	25	23	23	2,616	2,059	4,675	1.13	2.62
(%)	(49.2)	(17.5)	(9.8)	(8.3)	(7.6)	(3.1)	(1.7)	(1.0)	(0.9)	(0.9)	(100.0)				
総計	94,884	31,434	16,073	10,250	6,241	2,163	1,039	549	630	410	163,673	32,913	196,586	0.60	1.76
(%)	(58.0)	(19.2)	(9.8)	(6.3)	(3.8)	(1.3)	(0.6)	(0.3)	(0.4)	(0.3)	(100.0)				
平均	171.38	172.92	172.02	162.84	156.83	150.41	155.86	146.27	127.97	137.58	169.97	169.98	169.97		
標準偏差	153.91	162.55	167.77	169.57	158.92	169.80	238.88	180.29	114.04	149.80	159.10	170.39	160.97		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査