

表 3326 透析前CRP濃度(mg/dL) PTH値(pg/mL)別(施設血液透析患者 透析回数週3回) PTH測定法:whole-PTH

PTH値	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	691	263	155	96	82	34	11	5	17	11	1,365	134	1,499	1.04	3.57
(%)	(50.6)	(19.3)	(11.4)	(7.0)	(6.0)	(2.5)	(0.8)	(0.4)	(1.2)	(0.8)	(100.0)				
20～	961	313	166	108	69	33	12	9	20	9	1,700	148	1,848	0.88	3.26
(%)	(56.5)	(18.4)	(9.8)	(6.4)	(4.1)	(1.9)	(0.7)	(0.5)	(1.2)	(0.5)	(100.0)				
35～	826	266	167	108	62	23	8	8	15	6	1,489	124	1,613	0.77	2.32
(%)	(55.5)	(17.9)	(11.2)	(7.3)	(4.2)	(1.5)	(0.5)	(0.5)	(1.0)	(0.4)	(100.0)				
50～	1,244	427	224	156	85	34	15	13	22	16	2,236	223	2,459	0.89	3.55
(%)	(55.6)	(19.1)	(10.0)	(7.0)	(3.8)	(1.5)	(0.7)	(0.6)	(1.0)	(0.7)	(100.0)				
75～	1,073	402	201	118	78	24	14	10	8	9	1,937	211	2,148	0.75	3.02
(%)	(55.4)	(20.8)	(10.4)	(6.1)	(4.0)	(1.2)	(0.7)	(0.5)	(0.4)	(0.5)	(100.0)				
100～	910	331	177	109	55	21	6	10	11	7	1,637	150	1,787	0.71	2.47
(%)	(55.6)	(20.2)	(10.8)	(6.7)	(3.4)	(1.3)	(0.4)	(0.6)	(0.7)	(0.4)	(100.0)				
125～	618	203	122	54	32	15	12	7	7	3	1,073	114	1,187	0.67	1.80
(%)	(57.6)	(18.9)	(11.4)	(5.0)	(3.0)	(1.4)	(1.1)	(0.7)	(0.7)	(0.3)	(100.0)				
150～	15	8	3	2		1					29	8	37	0.48	1.07
(%)	(51.7)	(27.6)	(10.3)	(6.9)		(3.4)					(100.0)				
151～	453	194	67	56	23	9	5	1	8	2	818	116	934	0.61	1.67
(%)	(55.4)	(23.7)	(8.2)	(6.8)	(2.8)	(1.1)	(0.6)	(0.1)	(1.0)	(0.2)	(100.0)				
180～	326	132	58	25	23	5	4	4	4	6	587	79	666	0.84	2.96
(%)	(55.5)	(22.5)	(9.9)	(4.3)	(3.9)	(0.9)	(0.7)	(0.7)	(0.7)	(1.0)	(100.0)				
210～	200	105	32	41	15	7	2	3	3	2	410	62	472	0.95	4.72
(%)	(48.8)	(25.6)	(7.8)	(10.0)	(3.7)	(1.7)	(0.5)	(0.7)	(0.7)	(0.5)	(100.0)				
240～	280	82	43	30	15	1	3		3	2	459	67	526	0.58	1.89
(%)	(61.0)	(17.9)	(9.4)	(6.5)	(3.3)	(0.2)	(0.7)		(0.7)	(0.4)	(100.0)				
300～	107	42	11	13	4	1	1			3	182	23	205	0.75	2.97
(%)	(58.8)	(23.1)	(6.0)	(7.1)	(2.2)	(0.5)	(0.5)			(1.6)	(100.0)				
350～	69	18	17	4	7						115	20	135	0.41	0.68
(%)	(60.0)	(15.7)	(14.8)	(3.5)	(6.1)						(100.0)				
400～	68	30	14	12	3	2	1	1	1	3	135	17	152	1.26	4.32
(%)	(50.4)	(22.2)	(10.4)	(8.9)	(2.2)	(1.5)	(0.7)	(0.7)	(0.7)	(2.2)	(100.0)				
500～	74	28	12	10	10	2	1	1		2	140	26	166	0.95	2.91
(%)	(52.9)	(20.0)	(8.6)	(7.1)	(7.1)	(1.4)	(0.7)	(0.7)		(1.4)	(100.0)				
合計	7,915	2,844	1,469	942	563	212	95	72	119	81	14,312	1,522	15,834	0.80	2.97
(%)	(55.3)	(19.9)	(10.3)	(6.6)	(3.9)	(1.5)	(0.7)	(0.5)	(0.8)	(0.6)	(100.0)				
記載なし	61	19	9	6	6	4	2	1		1	109	329	438	0.97	2.20
(%)	(56.0)	(17.4)	(8.3)	(5.5)	(5.5)	(3.7)	(1.8)	(0.9)		(0.9)	(100.0)				
総計	7,976	2,863	1,478	948	569	216	97	73	119	82	14,421	1,851	16,272	0.81	2.97
(%)	(55.3)	(19.9)	(10.2)	(6.6)	(3.9)	(1.5)	(0.7)	(0.5)	(0.8)	(0.6)	(100.0)				
平均	104.51	107.17	100.43	104.06	100.03	86.25	98.46	97.10	80.76	137.14	104.05	121.15	105.70		
標準偏差	105.72	102.24	104.35	111.94	120.28	113.02	89.66	85.60	71.38	269.15	107.29	125.11	109.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査