

表 3331 透析前CRP濃度(mg/dL) 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回）

透析前収縮期血圧	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
60未満	15	5	1	3	4				1	2	31	4	35	2.42	5.60
(%)	(48.4)	(16.1)	(3.2)	(9.7)	(12.9)				(3.2)	(6.5)	(100.0)				
60～	74	52	24	30	20	12	4	3	6	4	229	42	271	1.76	3.49
(%)	(32.3)	(22.7)	(10.5)	(13.1)	(8.7)	(5.2)	(1.7)	(1.3)	(2.6)	(1.7)	(100.0)				
80～	828	378	275	195	152	53	40	22	26	20	1,989	388	2,377	1.31	3.14
(%)	(41.6)	(19.0)	(13.8)	(9.8)	(7.6)	(2.7)	(2.0)	(1.1)	(1.3)	(1.0)	(100.0)				
100～	5,406	2,167	1,233	899	668	258	164	85	98	75	11,053	2,102	13,155	1.01	2.70
(%)	(48.9)	(19.6)	(11.2)	(8.1)	(6.0)	(2.3)	(1.5)	(0.8)	(0.9)	(0.7)	(100.0)				
120～	19,379	6,756	3,531	2,348	1,519	537	269	164	195	131	34,829	6,747	41,576	0.71	2.14
(%)	(55.6)	(19.4)	(10.1)	(6.7)	(4.4)	(1.5)	(0.8)	(0.5)	(0.6)	(0.4)	(100.0)				
140～	31,834	10,274	5,030	3,309	1,885	662	316	168	213	141	53,832	10,371	64,203	0.59	1.89
(%)	(59.1)	(19.1)	(9.3)	(6.1)	(3.5)	(1.2)	(0.6)	(0.3)	(0.4)	(0.3)	(100.0)				
160～	23,357	7,561	3,738	2,315	1,344	469	210	117	139	80	39,330	7,513	46,843	0.56	1.70
(%)	(59.4)	(19.2)	(9.5)	(5.9)	(3.4)	(1.2)	(0.5)	(0.3)	(0.4)	(0.2)	(100.0)				
180～	8,791	3,001	1,501	925	540	181	92	49	33	26	15,139	2,863	18,002	0.55	1.52
(%)	(58.1)	(19.8)	(9.9)	(6.1)	(3.6)	(1.2)	(0.6)	(0.3)	(0.2)	(0.2)	(100.0)				
200～	2,274	762	403	212	120	41	21	4	12	6	3,855	660	4,515	0.51	1.35
(%)	(59.0)	(19.8)	(10.5)	(5.5)	(3.1)	(1.1)	(0.5)	(0.1)	(0.3)	(0.2)	(100.0)				
220～	449	131	71	52	29	9	5			2	748	146	894	0.53	1.40
(%)	(60.0)	(17.5)	(9.5)	(7.0)	(3.9)	(1.2)	(0.7)			(0.3)	(100.0)				
合計	92,407	31,087	15,807	10,288	6,281	2,222	1,121	612	723	487	161,035	30,836	191,871	0.64	1.96
(%)	(57.4)	(19.3)	(9.8)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.4)	(0.3)	(100.0)				
記載なし	13,122	4,291	2,340	1,484	973	360	161	82	119	73	23,005	14,095	37,100	0.66	1.83
(%)	(57.0)	(18.7)	(10.2)	(6.5)	(4.2)	(1.6)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
総計	105,529	35,378	18,147	11,772	7,254	2,582	1,282	694	842	560	184,040	44,931	228,971	0.64	1.94
(%)	(57.3)	(19.2)	(9.9)	(6.4)	(3.9)	(1.4)	(0.7)	(0.4)	(0.5)	(0.3)	(100.0)				
平均	152.78	151.71	150.90	149.58	147.51	146.73	144.76	143.49	141.97	140.53	151.72	151.23	151.64		
標準偏差	23.38	24.14	24.89	25.11	25.85	25.97	26.98	25.54	25.53	26.93	24.06	23.81	24.02		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査