

表 3337 透析前CRP濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	8,161	2,306	1,056	664	387	136	53	26	40	31	12,860	2,216	15,076	0.51	1.84
(%)	(63.5)	(17.9)	(8.2)	(5.2)	(3.0)	(1.1)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
慢性腎盂腎炎	225	67	33	22	6	5	3	1			362	60	422	0.44	1.01
(%)	(62.2)	(18.5)	(9.1)	(6.1)	(1.7)	(1.4)	(0.8)	(0.3)			(100.0)				
急速進行性糸球体腎炎	157	44	24	11	8	4	2	1			251	33	284	0.48	1.11
(%)	(62.5)	(17.5)	(9.6)	(4.4)	(3.2)	(1.6)	(0.8)	(0.4)			(100.0)				
妊娠腎/妊娠中毒症	200	50	23	14	8	3	1	2	1	1	303	56	359	0.51	1.66
(%)	(66.0)	(16.5)	(7.6)	(4.6)	(2.6)	(1.0)	(0.3)	(0.7)	(0.3)	(0.3)	(100.0)				
その他の分類不能の腎炎	105	41	10	8	5	1					170	25	195	0.34	0.68
(%)	(61.8)	(24.1)	(5.9)	(4.7)	(2.9)	(0.6)					(100.0)				
多発性嚢胞腎	711	262	119	79	49	15	4	4	6	4	1,253	211	1,464	0.59	1.60
(%)	(56.7)	(20.9)	(9.5)	(6.3)	(3.9)	(1.2)	(0.3)	(0.3)	(0.5)	(0.3)	(100.0)				
腎硬化症	1,435	495	269	163	81	32	13	10	5	5	2,508	378	2,886	0.56	1.59
(%)	(57.2)	(19.7)	(10.7)	(6.5)	(3.2)	(1.3)	(0.5)	(0.4)	(0.2)	(0.2)	(100.0)				
悪性高血圧	154	70	24	12	6	2		1	1	1	271	41	312	0.47	1.42
(%)	(56.8)	(25.8)	(8.9)	(4.4)	(2.2)	(0.7)		(0.4)	(0.4)	(0.4)	(100.0)				
糖尿病性腎症	6,610	2,240	1,133	763	412	144	65	41	37	35	11,480	1,693	13,173	0.61	1.82
(%)	(57.6)	(19.5)	(9.9)	(6.6)	(3.6)	(1.3)	(0.6)	(0.4)	(0.3)	(0.3)	(100.0)				
SLE腎炎	193	56	30	23	11	6	5				324	39	363	0.55	1.15
(%)	(59.6)	(17.3)	(9.3)	(7.1)	(3.4)	(1.9)	(1.5)				(100.0)				
アミロイド腎	32	11	5	3	4						55	4	59	0.51	0.94
(%)	(58.2)	(20.0)	(9.1)	(5.5)	(7.3)						(100.0)				
痛風腎	71	26	14	11	5	2	1				130	24	154	0.51	0.95
(%)	(54.6)	(20.0)	(10.8)	(8.5)	(3.8)	(1.5)	(0.8)				(100.0)				
先天性代謝異常による腎不全	24	6	1	1	2	2					36	8	44	0.53	1.14
(%)	(66.7)	(16.7)	(2.8)	(2.8)	(5.6)	(5.6)					(100.0)				
腎・尿路結核	12	1	3	1		1					18	4	22	0.50	0.98
(%)	(66.7)	(5.6)	(16.7)	(5.6)		(5.6)					(100.0)				
腎・尿路結石	33	14	4	5	3						59	10	69	0.45	0.77
(%)	(55.9)	(23.7)	(6.8)	(8.5)	(5.1)						(100.0)				
腎・尿路腫瘍	35	15	12	5	4					2	73	12	85	1.30	5.16
(%)	(47.9)	(20.5)	(16.4)	(6.8)	(5.5)					(2.7)	(100.0)				
閉塞性尿路障害	40	10	8	2	2	2			1		65	9	74	0.65	1.72
(%)	(61.5)	(15.4)	(12.3)	(3.1)	(3.1)	(3.1)			(1.5)		(100.0)				
骨髄腫	8	4	4	1	1						18	1	19	0.47	0.58
(%)	(44.4)	(22.2)	(22.2)	(5.6)	(5.6)						(100.0)				
腎形成不全	56	18	4	4	4						86	15	101	0.32	0.60
(%)	(65.1)	(20.9)	(4.7)	(4.7)	(4.7)						(100.0)				
不明	1,804	527	293	140	94	34	16	7	13	3	2,931	549	3,480	0.52	1.64
(%)	(61.5)	(18.0)	(10.0)	(4.8)	(3.2)	(1.2)	(0.5)	(0.2)	(0.4)	(0.1)	(100.0)				
再導入	207	79	30	18	17	4	1				356	60	416	0.43	0.82
(%)	(58.1)	(22.2)	(8.4)	(5.1)	(4.8)	(1.1)	(0.3)				(100.0)				
その他	431	135	74	53	25	14	7	1	4	2	746	143	889	0.65	1.63
(%)	(57.8)	(18.1)	(9.9)	(7.1)	(3.4)	(1.9)	(0.9)	(0.1)	(0.5)	(0.3)	(100.0)				
合計	20,704	6,477	3,173	2,003	1,134	407	171	94	108	84	34,355	5,591	39,946	0.55	1.76
(%)	(60.3)	(18.9)	(9.2)	(5.8)	(3.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
記載なし															
(%)															
総計	20,704	6,477	3,173	2,003	1,134	407	171	94	108	84	34,355	5,591	39,946	0.55	1.76
(%)	(60.3)	(18.9)	(9.2)	(5.8)	(3.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査