

表 3342 透析前CRP濃度(mg/dL) PTH値(pg/mL)別(血液透析濾過患者 透析回数週3回) PTH測定法:intact-PTH

PTH値	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	839	301	148	115	52	24	13	6	10	6	1,514	169	1,683	0.70	1.90
(%)	(55.4)	(19.9)	(9.8)	(7.6)	(3.4)	(1.6)	(0.9)	(0.4)	(0.7)	(0.4)	(100.0)				
20～	1,151	343	190	118	76	25	14	2	6	7	1,932	263	2,195	0.61	2.06
(%)	(59.6)	(17.8)	(9.8)	(6.1)	(3.9)	(1.3)	(0.7)	(0.1)	(0.3)	(0.4)	(100.0)				
40～	1,270	391	213	146	91	37	11	6	5	5	2,175	262	2,437	0.60	1.69
(%)	(58.4)	(18.0)	(9.8)	(6.7)	(4.2)	(1.7)	(0.5)	(0.3)	(0.2)	(0.2)	(100.0)				
60～	1,377	436	203	132	89	23	16	9	6	6	2,297	295	2,592	0.57	1.57
(%)	(59.9)	(19.0)	(8.8)	(5.7)	(3.9)	(1.0)	(0.7)	(0.4)	(0.3)	(0.3)	(100.0)				
80～	1,504	487	249	162	88	23	11	8	4	9	2,545	301	2,846	0.56	2.17
(%)	(59.1)	(19.1)	(9.8)	(6.4)	(3.5)	(0.9)	(0.4)	(0.3)	(0.2)	(0.4)	(100.0)				
100～	1,504	464	224	151	73	33	14	12	14	6	2,495	300	2,795	0.59	1.68
(%)	(60.3)	(18.6)	(9.0)	(6.1)	(2.9)	(1.3)	(0.6)	(0.5)	(0.6)	(0.2)	(100.0)				
120～	1,390	443	175	129	73	35	10	6	6		2,267	314	2,581	0.49	1.19
(%)	(61.3)	(19.5)	(7.7)	(5.7)	(3.2)	(1.5)	(0.4)	(0.3)	(0.3)		(100.0)				
140～	1,269	392	187	114	60	23	9	4	12	3	2,073	250	2,323	0.51	1.42
(%)	(61.2)	(18.9)	(9.0)	(5.5)	(2.9)	(1.1)	(0.4)	(0.2)	(0.6)	(0.1)	(100.0)				
160～	1,199	343	176	105	45	23	2	7	3	4	1,907	243	2,150	0.48	1.68
(%)	(62.9)	(18.0)	(9.2)	(5.5)	(2.4)	(1.2)	(0.1)	(0.4)	(0.2)	(0.2)	(100.0)				
180～	2,780	853	401	227	123	48	12	10	5	6	4,465	540	5,005	0.45	1.40
(%)	(62.3)	(19.1)	(9.0)	(5.1)	(2.8)	(1.1)	(0.3)	(0.2)	(0.1)	(0.1)	(100.0)				
240～	41	11	6	7	5						70	7	77	0.50	0.72
(%)	(58.6)	(15.7)	(8.6)	(10.0)	(7.1)						(100.0)				
241～	1,653	496	250	118	92	25	13	2	7	1	2,657	298	2,955	0.45	1.16
(%)	(62.2)	(18.7)	(9.4)	(4.4)	(3.5)	(0.9)	(0.5)	(0.1)	(0.3)	(0.0)	(100.0)				
300～	945	267	129	94	44	12	10	2	4	5	1,512	202	1,714	0.52	1.51
(%)	(62.5)	(17.7)	(8.5)	(6.2)	(2.9)	(0.8)	(0.7)	(0.1)	(0.3)	(0.3)	(100.0)				
360～	553	183	98	43	21	6	3	4	3	2	916	126	1,042	0.51	1.74
(%)	(60.4)	(20.0)	(10.7)	(4.7)	(2.3)	(0.7)	(0.3)	(0.4)	(0.3)	(0.2)	(100.0)				
420～	323	102	46	29	16	6	6	1		1	530	61	591	0.52	1.47
(%)	(60.9)	(19.2)	(8.7)	(5.5)	(3.0)	(1.1)	(1.1)	(0.2)		(0.2)	(100.0)				
480～	63	32	18	10	3	1					127	19	146	0.42	0.64
(%)	(49.6)	(25.2)	(14.2)	(7.9)	(2.4)	(0.8)					(100.0)				
500～	121	43	27	15	2	3				1	212	25	237	0.46	1.40
(%)	(57.1)	(20.3)	(12.7)	(7.1)	(0.9)	(1.4)				(0.5)	(100.0)				
540～	133	40	20	9	8	1					211	41	252	0.36	0.67
(%)	(63.0)	(19.0)	(9.5)	(4.3)	(3.8)	(0.5)					(100.0)				
600～	211	69	42	19	15	4	1	1			362	61	423	0.46	0.96
(%)	(58.3)	(19.1)	(11.6)	(5.2)	(4.1)	(1.1)	(0.3)	(0.3)			(100.0)				
800～	165	67	26	18	13	3	4				296	54	350	0.52	1.04
(%)	(55.7)	(22.6)	(8.8)	(6.1)	(4.4)	(1.0)	(1.4)				(100.0)				
合計	18,491	5,763	2,828	1,761	989	355	149	80	85	62	30,563	3,831	34,394	0.53	1.60
(%)	(60.5)	(18.9)	(9.3)	(5.8)	(3.2)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
記載なし	230	67	38	29	30	11	8	2	3		418	316	734	0.83	1.73
(%)	(55.0)	(16.0)	(9.1)	(6.9)	(7.2)	(2.6)	(1.9)	(0.5)	(0.7)		(100.0)				
総計	18,721	5,830	2,866	1,790	1,019	366	157	82	88	62	30,981	4,147	35,128	0.53	1.60
(%)	(60.4)	(18.8)	(9.3)	(5.8)	(3.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
平均	177.01	179.18	178.47	169.74	170.39	159.21	169.09	144.54	132.98	133.60	176.38	184.27	177.26		
標準偏差	164.54	174.07	164.13	184.64	180.36	149.56	178.57	110.75	95.51	117.41	167.69	182.83	169.45		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査