

表 3347 透析前CRP濃度(mg/dL) Non-HDL-C濃度(mg/dL)別(血液透析濾過患者 透析回数週3回)

Non-HDL-C濃度	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
50未満	298	66	54	30	19	10	8	3	1	1	490	39	529	0.68	1.67
(%)	(60.8)	(13.5)	(11.0)	(6.1)	(3.9)	(2.0)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
50～	1,309	379	197	128	100	34	14	8	6	3	2,178	225	2,403	0.58	1.43
(%)	(60.1)	(17.4)	(9.0)	(5.9)	(4.6)	(1.6)	(0.6)	(0.4)	(0.3)	(0.1)	(100.0)				
70～	2,864	832	414	268	153	64	26	15	14	7	4,657	481	5,138	0.54	1.55
(%)	(61.5)	(17.9)	(8.9)	(5.8)	(3.3)	(1.4)	(0.6)	(0.3)	(0.3)	(0.2)	(100.0)				
90～	3,510	1,017	492	318	168	65	32	15	17	16	5,650	615	6,265	0.54	1.72
(%)	(62.1)	(18.0)	(8.7)	(5.6)	(3.0)	(1.2)	(0.6)	(0.3)	(0.3)	(0.3)	(100.0)				
110～	2,860	902	407	237	148	37	19	5	15	10	4,640	522	5,162	0.49	1.48
(%)	(61.6)	(19.4)	(8.8)	(5.1)	(3.2)	(0.8)	(0.4)	(0.1)	(0.3)	(0.2)	(100.0)				
130～	1,746	560	286	146	82	25	7	5	4	2	2,863	286	3,149	0.42	1.03
(%)	(61.0)	(19.6)	(10.0)	(5.1)	(2.9)	(0.9)	(0.2)	(0.2)	(0.1)	(0.1)	(100.0)				
150～	839	287	156	94	39	12	3	1	2	3	1,436	161	1,597	0.47	1.32
(%)	(58.4)	(20.0)	(10.9)	(6.5)	(2.7)	(0.8)	(0.2)	(0.1)	(0.1)	(0.2)	(100.0)				
170～	352	126	70	45	10	7		1	1		612	70	682	0.43	0.94
(%)	(57.5)	(20.6)	(11.4)	(7.4)	(1.6)	(1.1)		(0.2)	(0.2)		(100.0)				
190～	122	50	28	21	10				2		233	29	262	0.56	1.32
(%)	(52.4)	(21.5)	(12.0)	(9.0)	(4.3)				(0.9)		(100.0)				
210～	55	36	22	9	3	2	1	1			129	12	141	0.60	1.23
(%)	(42.6)	(27.9)	(17.1)	(7.0)	(2.3)	(1.6)	(0.8)	(0.8)			(100.0)				
合計	13,955	4,255	2,126	1,296	732	256	110	54	62	42	22,888	2,440	25,328	0.51	1.48
(%)	(61.0)	(18.6)	(9.3)	(5.7)	(3.2)	(1.1)	(0.5)	(0.2)	(0.3)	(0.2)	(100.0)				
記載なし	6,749	2,222	1,047	707	402	151	61	40	46	42	11,467	3,151	14,618	0.63	2.21
(%)	(58.9)	(19.4)	(9.1)	(6.2)	(3.5)	(1.3)	(0.5)	(0.3)	(0.4)	(0.4)	(100.0)				
総計	20,704	6,477	3,173	2,003	1,134	407	171	94	108	84	34,355	5,591	39,946	0.55	1.76
(%)	(60.3)	(18.9)	(9.2)	(5.8)	(3.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
平均	106.40	109.35	109.49	107.82	102.63	99.48	95.18	93.70	103.56	103.05	107.02	108.04	107.12		
標準偏差	32.50	33.85	36.15	35.95	33.33	34.01	32.15	36.90	31.83	25.42	33.41	33.03	33.37		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査