

表 3349 透析前CRP濃度(mg/dL) 透析前拡張期血圧(mmHg)別（血液透析濾過患者 透析回数週3回）

透析前拡張 期血圧	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	2	2		1	1						6	1	7	0.93	1.26
(%)	(33.3)	(33.3)		(16.7)	(16.7)						(100.0)				
20～	40	27	11	15	6	4		1		1	105	16	121	1.02	2.17
(%)	(38.1)	(25.7)	(10.5)	(14.3)	(5.7)	(3.8)		(1.0)		(1.0)	(100.0)				
40～	1,131	487	278	178	117	47	25	18	16	13	2,310	370	2,680	0.88	2.29
(%)	(49.0)	(21.1)	(12.0)	(7.7)	(5.1)	(2.0)	(1.1)	(0.8)	(0.7)	(0.6)	(100.0)				
60～	7,786	2,460	1,252	801	475	180	67	39	43	31	13,134	1,838	14,972	0.58	1.83
(%)	(59.3)	(18.7)	(9.5)	(6.1)	(3.6)	(1.4)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
80～	4,962	1,503	668	441	221	72	29	19	21	18	7,954	1,049	9,003	0.49	1.50
(%)	(62.4)	(18.9)	(8.4)	(5.5)	(2.8)	(0.9)	(0.4)	(0.2)	(0.3)	(0.2)	(100.0)				
90～	3,041	845	394	242	120	39	27	4	4	7	4,723	594	5,317	0.45	1.76
(%)	(64.4)	(17.9)	(8.3)	(5.1)	(2.5)	(0.8)	(0.6)	(0.1)	(0.1)	(0.1)	(100.0)				
100～	1,524	431	225	119	68	26	3	3	7	1	2,407	294	2,701	0.43	1.13
(%)	(63.3)	(17.9)	(9.3)	(4.9)	(2.8)	(1.1)	(0.1)	(0.1)	(0.3)	(0.0)	(100.0)				
120～	89	30	20	6	4	2					151	18	169	0.39	0.72
(%)	(58.9)	(19.9)	(13.2)	(4.0)	(2.6)	(1.3)					(100.0)				
140～	5	1	1	1		1					9	3	12	0.85	1.74
(%)	(55.6)	(11.1)	(11.1)	(11.1)		(11.1)					(100.0)				
合計	18,580	5,786	2,849	1,804	1,012	371	151	84	91	71	30,799	4,183	34,982	0.55	1.73
(%)	(60.3)	(18.8)	(9.3)	(5.9)	(3.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
記載なし	2,124	691	324	199	122	36	20	10	17	13	3,556	1,408	4,964	0.60	1.97
(%)	(59.7)	(19.4)	(9.1)	(5.6)	(3.4)	(1.0)	(0.6)	(0.3)	(0.5)	(0.4)	(100.0)				
総計	20,704	6,477	3,173	2,003	1,134	407	171	94	108	84	34,355	5,591	39,946	0.55	1.76
(%)	(60.3)	(18.9)	(9.2)	(5.8)	(3.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(100.0)				
平均	80.17	78.62	78.36	77.57	76.03	75.54	74.21	71.14	73.66	72.49	79.28	78.39	79.17		
標準偏差	14.26	14.89	15.36	15.16	15.70	16.54	14.91	15.02	15.65	15.18	14.69	14.80	14.70		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査