

表 3354 CRP濃度(mg/dL) 原疾患別（腹膜透析患者）

原疾患	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,120	307	147	102	49	28	9	4	13	3	1,782	1,277	3,059	0.56	1.63
(%)	(62.9)	(17.2)	(8.2)	(5.7)	(2.7)	(1.6)	(0.5)	(0.2)	(0.7)	(0.2)	(100.0)				
慢性腎盂腎炎	37	9	2	2	2	2		1	1		56	37	93	0.82	2.14
(%)	(66.1)	(16.1)	(3.6)	(3.6)	(3.6)	(3.6)		(1.8)	(1.8)		(100.0)				
急速進行性糸球体腎炎	30	8	3	2	2	1					46	28	74	0.41	0.89
(%)	(65.2)	(17.4)	(6.5)	(4.3)	(4.3)	(2.2)					(100.0)				
妊娠腎/妊娠中毒症	8	6	4	1	1	1					21	11	32	0.65	1.07
(%)	(38.1)	(28.6)	(19.0)	(4.8)	(4.8)	(4.8)					(100.0)				
その他の分類不能の腎炎	40	7	2	2	2	1					54	31	85	0.34	0.79
(%)	(74.1)	(13.0)	(3.7)	(3.7)	(3.7)	(1.9)					(100.0)				
多発性嚢胞腎	76	22	8	9	8	1	1		2	1	128	115	243	0.83	2.39
(%)	(59.4)	(17.2)	(6.3)	(7.0)	(6.3)	(0.8)	(0.8)		(1.6)	(0.8)	(100.0)				
腎硬化症	370	139	52	40	29	9	6	6	6		657	426	1,083	0.68	1.63
(%)	(56.3)	(21.2)	(7.9)	(6.1)	(4.4)	(1.4)	(0.9)	(0.9)	(0.9)		(100.0)				
悪性高血圧	35	10	12	4	2		1				64	43	107	0.48	0.98
(%)	(54.7)	(15.6)	(18.8)	(6.3)	(3.1)		(1.6)				(100.0)				
糖尿病性腎症	861	344	154	97	75	27	16	8	9	2	1,593	1,173	2,766	0.69	1.92
(%)	(54.0)	(21.6)	(9.7)	(6.1)	(4.7)	(1.7)	(1.0)	(0.5)	(0.6)	(0.1)	(100.0)				
SLE腎炎	18	9	5	3	2			2			39	23	62	0.88	2.00
(%)	(46.2)	(23.1)	(12.8)	(7.7)	(5.1)			(5.1)			(100.0)				
アミロイド腎	2				1						3	6	9	1.29	1.99
(%)	(66.7)				(33.3)						(100.0)				
痛風腎	16	4	1								21	12	33	0.15	0.22
(%)	(76.2)	(19.0)	(4.8)								(100.0)				
先天性代謝異常による腎不全	9	3			1						13	5	18	0.26	0.56
(%)	(69.2)	(23.1)			(7.7)						(100.0)				
腎・尿路結核	1										1		1		
(%)	(100.0)										(100.0)				
腎・尿路結石	6			1		1					8	8	16	0.89	1.89
(%)	(75.0)			(12.5)		(12.5)					(100.0)				
腎・尿路腫瘍	3	2		2	1						8	6	14	0.87	0.94
(%)	(37.5)	(25.0)		(25.0)	(12.5)						(100.0)				
閉塞性尿路障害	10	2		1		1					14	8	22	0.54	1.32
(%)	(71.4)	(14.3)		(7.1)		(7.1)					(100.0)				
骨髄腫	3			1		1					5	3	8	1.20	1.87
(%)	(60.0)			(20.0)		(20.0)					(100.0)				
腎形成不全	37	4	4	1	2						48	28	76	0.29	0.70
(%)	(77.1)	(8.3)	(8.3)	(2.1)	(4.2)						(100.0)				
不明	275	90	38	31	20	7	7	1	3	1	473	367	840	0.66	1.62
(%)	(58.1)	(19.0)	(8.0)	(6.6)	(4.2)	(1.5)	(1.5)	(0.2)	(0.6)	(0.2)	(100.0)				
再導入	21	8	3			1					33	18	51	0.34	0.94
(%)	(63.6)	(24.2)	(9.1)			(3.0)					(100.0)				
その他	89	21	14	9	5	2		2			142	107	249	0.53	1.25
(%)	(62.7)	(14.8)	(9.9)	(6.3)	(3.5)	(1.4)		(1.4)			(100.0)				
合計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
記載なし															
(%)															
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査