

表 3356 CRP濃度(mg/dL) 体重(kg)別 (腹膜透析患者)

体重	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	28	2	3		1						34	4	38	0.17	0.37
(%)	(82.4)	(5.9)	(8.8)		(2.9)						(100.0)				
20～	10	4	4		2						20	3	23	0.47	0.75
(%)	(50.0)	(20.0)	(20.0)		(10.0)						(100.0)				
30～	81	26	15	7	9	1	3	1	2	1	146	26	172	0.94	2.46
(%)	(55.5)	(17.8)	(10.3)	(4.8)	(6.2)	(0.7)	(2.1)	(0.7)	(1.4)	(0.7)	(100.0)				
40～	523	115	51	36	21	10	6	6	5		773	131	904	0.54	1.51
(%)	(67.7)	(14.9)	(6.6)	(4.7)	(2.7)	(1.3)	(0.8)	(0.8)	(0.6)		(100.0)				
50～	830	235	105	81	56	24	11	7	5	2	1,356	235	1,591	0.61	1.61
(%)	(61.2)	(17.3)	(7.7)	(6.0)	(4.1)	(1.8)	(0.8)	(0.5)	(0.4)	(0.1)	(100.0)				
60～	806	269	118	84	50	21	7	3	10	1	1,369	247	1,616	0.57	1.48
(%)	(58.9)	(19.6)	(8.6)	(6.1)	(3.7)	(1.5)	(0.5)	(0.2)	(0.7)	(0.1)	(100.0)				
70～	382	162	64	38	21	6	4	1	6	1	685	126	811	0.60	2.04
(%)	(55.8)	(23.6)	(9.3)	(5.5)	(3.1)	(0.9)	(0.6)	(0.1)	(0.9)	(0.1)	(100.0)				
80～	130	59	29	17	7	4		2	2		250	48	298	0.62	1.55
(%)	(52.0)	(23.6)	(11.6)	(6.8)	(2.8)	(1.6)		(0.8)	(0.8)		(100.0)				
90～	31	30	10	4	7	5					87	17	104	0.77	1.24
(%)	(35.6)	(34.5)	(11.5)	(4.6)	(8.0)	(5.7)					(100.0)				
100～	4	3	2	2	4		1				16	4	20	1.38	1.64
(%)	(25.0)	(18.8)	(12.5)	(12.5)	(25.0)		(6.3)				(100.0)				
110～	3	3			1	1					8	2	10	1.25	1.96
(%)	(37.5)	(37.5)			(12.5)	(12.5)					(100.0)				
120～	5	2			1	1					9		9	1.06	1.89
(%)	(55.6)	(22.2)			(11.1)	(11.1)					(100.0)				
合計	2,833	910	401	269	180	73	32	20	30	5	4,753	843	5,596	0.60	1.65
(%)	(59.6)	(19.1)	(8.4)	(5.7)	(3.8)	(1.5)	(0.7)	(0.4)	(0.6)	(0.1)	(100.0)				
記載なし	234	85	48	39	22	10	8	4	4	2	456	2,889	3,345	0.91	2.26
(%)	(51.3)	(18.6)	(10.5)	(8.6)	(4.8)	(2.2)	(1.8)	(0.9)	(0.9)	(0.4)	(100.0)				
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
平均	59.51	63.03	61.80	61.73	62.17	63.72	57.70	57.35	61.21	54.48	60.65	61.27	60.74		
標準偏差	13.55	14.12	14.17	12.56	17.11	16.73	13.92	13.02	13.90	12.82	13.94	13.65	13.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査