

表 3361 CRP濃度(mg/dL) ヘモグロビン濃度(g/dL)別 (腹膜透析患者)

ヘモグロビン 濃度	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
5未満	2	2		2							6	1	7	0.62	0.70
(%)	(33.3)	(33.3)		(33.3)							(100.0)				
5～	6	1	4		2						13	1	14	0.76	0.98
(%)	(46.2)	(7.7)	(30.8)		(15.4)						(100.0)				
6～	6	6	3	5	2	2				1	25	3	28	1.52	2.50
(%)	(24.0)	(24.0)	(12.0)	(20.0)	(8.0)	(8.0)				(4.0)	(100.0)				
7～	51	20	11	6	13	4	4	1	1	1	112	12	124	1.48	4.08
(%)	(45.5)	(17.9)	(9.8)	(5.4)	(11.6)	(3.6)	(3.6)	(0.9)	(0.9)	(0.9)	(100.0)				
8～	138	59	24	24	22	14	7	4	5		297	47	344	1.17	2.22
(%)	(46.5)	(19.9)	(8.1)	(8.1)	(7.4)	(4.7)	(2.4)	(1.3)	(1.7)		(100.0)				
9～	431	153	61	70	42	16	10	3	5	2	793	110	903	0.75	1.70
(%)	(54.4)	(19.3)	(7.7)	(8.8)	(5.3)	(2.0)	(1.3)	(0.4)	(0.6)	(0.3)	(100.0)				
10～	843	289	150	81	52	27	14	7	8	1	1,472	234	1,706	0.62	1.61
(%)	(57.3)	(19.6)	(10.2)	(5.5)	(3.5)	(1.8)	(1.0)	(0.5)	(0.5)	(0.1)	(100.0)				
11～	952	268	104	72	41	13	2	4	9	1	1,466	243	1,709	0.46	1.35
(%)	(64.9)	(18.3)	(7.1)	(4.9)	(2.8)	(0.9)	(0.1)	(0.3)	(0.6)	(0.1)	(100.0)				
12～	439	135	58	29	17	6	1	2	4	1	692	104	796	0.47	1.43
(%)	(63.4)	(19.5)	(8.4)	(4.2)	(2.5)	(0.9)	(0.1)	(0.3)	(0.6)	(0.1)	(100.0)				
13～	125	41	22	7	8			1	1	1	206	31	237	0.57	2.27
(%)	(60.7)	(19.9)	(10.7)	(3.4)	(3.9)			(0.5)	(0.5)	(0.5)	(100.0)				
14～	38	14	8	7	1		1	2			71	8	79	0.70	1.61
(%)	(53.5)	(19.7)	(11.3)	(9.9)	(1.4)		(1.4)	(2.8)			(100.0)				
合計	3,031	988	445	303	200	82	39	24	34	7	5,153	794	5,947	0.63	1.72
(%)	(58.8)	(19.2)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
記載なし	36	7	4	5	2	1	1				56	2,938	2,994	0.59	1.27
(%)	(64.3)	(12.5)	(7.1)	(8.9)	(3.6)	(1.8)	(1.8)				(100.0)				
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
平均	10.92	10.80	10.80	10.49	10.24	9.99	9.66	10.74	10.36	10.57	10.81	10.85	10.81		
標準偏差	1.32	1.46	1.53	1.62	1.61	1.38	1.40	2.10	1.57	1.82	1.42	1.37	1.42		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査