

表 3363 CRP濃度(mg/dL) HDL-C濃度(mg/dL)別(腹膜透析患者)

HDL-C濃度	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	2	4	1		1						8		8	0.55	0.71
(%)	(25.0)	(50.0)	(12.5)		(12.5)						(100.0)				
20～	95	66	42	27	16	7	2	1	5		261	27	288	0.98	1.95
(%)	(36.4)	(25.3)	(16.1)	(10.3)	(6.1)	(2.7)	(0.8)	(0.4)	(1.9)		(100.0)				
30～	426	199	85	59	43	17	3	2	3	5	842	91	933	0.76	2.38
(%)	(50.6)	(23.6)	(10.1)	(7.0)	(5.1)	(2.0)	(0.4)	(0.2)	(0.4)	(0.6)	(100.0)				
40～	504	194	81	59	31	12	12	2	4		899	91	990	0.58	1.35
(%)	(56.1)	(21.6)	(9.0)	(6.6)	(3.4)	(1.3)	(1.3)	(0.2)	(0.4)		(100.0)				
50～	510	130	58	30	19	12		3	2		764	75	839	0.41	1.07
(%)	(66.8)	(17.0)	(7.6)	(3.9)	(2.5)	(1.6)		(0.4)	(0.3)		(100.0)				
60～	338	49	37	14	11	4	2	2			457	59	516	0.36	0.98
(%)	(74.0)	(10.7)	(8.1)	(3.1)	(2.4)	(0.9)	(0.4)	(0.4)			(100.0)				
70～	185	32	9	5	4						235	24	259	0.19	0.43
(%)	(78.7)	(13.6)	(3.8)	(2.1)	(1.7)						(100.0)				
80～	97	16	4	7	5	2	1		1		133	8	141	0.51	1.46
(%)	(72.9)	(12.0)	(3.0)	(5.3)	(3.8)	(1.5)	(0.8)		(0.8)		(100.0)				
90～	46	6	2	2		2					58	3	61	0.32	0.91
(%)	(79.3)	(10.3)	(3.4)	(3.4)		(3.4)					(100.0)				
100～	38	9	1		1		1	1			51	6	57	0.50	1.63
(%)	(74.5)	(17.6)	(2.0)		(2.0)		(2.0)	(2.0)			(100.0)				
合計	2,241	705	320	203	131	56	21	11	15	5	3,708	384	4,092	0.56	1.59
(%)	(60.4)	(19.0)	(8.6)	(5.5)	(3.5)	(1.5)	(0.6)	(0.3)	(0.4)	(0.1)	(100.0)				
記載なし	826	290	129	105	71	27	19	13	19	2	1,501	3,348	4,849	0.81	1.97
(%)	(55.0)	(19.3)	(8.6)	(7.0)	(4.7)	(1.8)	(1.3)	(0.9)	(1.3)	(0.1)	(100.0)				
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
平均	53.56	46.84	45.28	44.27	44.48	45.86	48.81	54.91	39.47	34.40	50.52	50.04	50.47		
標準偏差	18.04	16.91	14.64	15.28	15.64	15.99	21.81	28.78	17.13	3.21	17.77	17.31	17.72		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査