

表 3364 CRP濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
50未満	12	3	1	1							17	1	18	0.23	0.31
(%)	(70.6)	(17.6)	(5.9)	(5.9)							(100.0)				
50～	57	18	8	7	7	3	2	1		1	104	10	114	1.18	4.08
(%)	(54.8)	(17.3)	(7.7)	(6.7)	(6.7)	(2.9)	(1.9)	(1.0)		(1.0)	(100.0)				
70～	194	63	21	13	7	9	2	1	3	1	314	37	351	0.71	2.33
(%)	(61.8)	(20.1)	(6.7)	(4.1)	(2.2)	(2.9)	(0.6)	(0.3)	(1.0)	(0.3)	(100.0)				
90～	370	116	68	20	19	11	3	1	3	1	612	48	660	0.52	1.38
(%)	(60.5)	(19.0)	(11.1)	(3.3)	(3.1)	(1.8)	(0.5)	(0.2)	(0.5)	(0.2)	(100.0)				
110～	443	140	55	43	26	9	5	3	2		726	71	797	0.52	1.24
(%)	(61.0)	(19.3)	(7.6)	(5.9)	(3.6)	(1.2)	(0.7)	(0.4)	(0.3)		(100.0)				
130～	391	112	45	35	21	5	3	1	3		616	46	662	0.47	1.20
(%)	(63.5)	(18.2)	(7.3)	(5.7)	(3.4)	(0.8)	(0.5)	(0.2)	(0.5)		(100.0)				
150～	241	76	36	25	14	4	2	1			399	37	436	0.44	0.92
(%)	(60.4)	(19.0)	(9.0)	(6.3)	(3.5)	(1.0)	(0.5)	(0.3)			(100.0)				
170～	134	49	27	18	12	2		1	3		246	17	263	0.63	1.53
(%)	(54.5)	(19.9)	(11.0)	(7.3)	(4.9)	(0.8)		(0.4)	(1.2)		(100.0)				
190～	70	17	10	5	3	2	1				108	12	120	0.49	1.11
(%)	(64.8)	(15.7)	(9.3)	(4.6)	(2.8)	(1.9)	(0.9)				(100.0)				
210～	39	20	12	5	7	1		1			85	5	90	0.65	1.28
(%)	(45.9)	(23.5)	(14.1)	(5.9)	(8.2)	(1.2)		(1.2)			(100.0)				
合計	1,951	614	283	172	116	46	18	10	14	3	3,227	284	3,511	0.55	1.56
(%)	(60.5)	(19.0)	(8.8)	(5.3)	(3.6)	(1.4)	(0.6)	(0.3)	(0.4)	(0.1)	(100.0)				
記載なし	1,116	381	166	136	86	37	22	14	20	4	1,982	3,448	5,430	0.76	1.93
(%)	(56.3)	(19.2)	(8.4)	(6.9)	(4.3)	(1.9)	(1.1)	(0.7)	(1.0)	(0.2)	(100.0)				
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
平均	127.67	129.33	131.86	133.40	135.20	116.91	117.56	139.20	126.43	80.67	128.70	125.82	128.47		
標準偏差	36.73	37.98	41.96	38.25	44.76	38.19	37.64	70.63	36.94	27.74	38.08	37.25	38.01		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査