

表 3366 CRP濃度(mg/dL) 拡張期血圧(mmHg)別（腹膜透析患者）

拡張期血圧	0.2未満	0.2～	0.5～	1.0～	2.0～	4.0～	6.0～	8.0～	10.0～	15.0～	合計	記載なし	総計	平均	標準偏差
20未満	1										1		1	0.18	
(%)	(100.0)										(100.0)				
20～	3	4		2	1	2					12	3	15	1.22	1.65
(%)	(25.0)	(33.3)		(16.7)	(8.3)	(16.7)					(100.0)				
40～	150	62	43	26	17	12	5	2	2	2	321	54	375	1.02	2.82
(%)	(46.7)	(19.3)	(13.4)	(8.1)	(5.3)	(3.7)	(1.6)	(0.6)	(0.6)	(0.6)	(100.0)				
60～	1,057	333	146	111	70	32	16	3	13	4	1,785	280	2,065	0.65	1.81
(%)	(59.2)	(18.7)	(8.2)	(6.2)	(3.9)	(1.8)	(0.9)	(0.2)	(0.7)	(0.2)	(100.0)				
80～	672	226	88	63	40	10	4	8	8		1,119	177	1,296	0.56	1.50
(%)	(60.1)	(20.2)	(7.9)	(5.6)	(3.6)	(0.9)	(0.4)	(0.7)	(0.7)		(100.0)				
90～	402	112	56	27	17	7	3		2		626	88	714	0.42	1.08
(%)	(64.2)	(17.9)	(8.9)	(4.3)	(2.7)	(1.1)	(0.5)		(0.3)		(100.0)				
100～	191	58	17	11	11	3		1	1		293	54	347	0.42	1.13
(%)	(65.2)	(19.8)	(5.8)	(3.8)	(3.8)	(1.0)		(0.3)	(0.3)		(100.0)				
120～	9	3	1	1							14	4	18	0.28	0.37
(%)	(64.3)	(21.4)	(7.1)	(7.1)							(100.0)				
140～						1					1		1	4.76	
(%)						(100.0)					(100.0)				
合計	2,485	798	351	241	156	67	28	14	26	6	4,172	660	4,832	0.61	1.71
(%)	(59.6)	(19.1)	(8.4)	(5.8)	(3.7)	(1.6)	(0.7)	(0.3)	(0.6)	(0.1)	(100.0)				
記載なし	582	197	98	67	46	16	12	10	8	1	1,037	3,072	4,109	0.72	1.72
(%)	(56.1)	(19.0)	(9.5)	(6.5)	(4.4)	(1.5)	(1.2)	(1.0)	(0.8)	(0.1)	(100.0)				
総計	3,067	995	449	308	202	83	40	24	34	7	5,209	3,732	8,941	0.63	1.71
(%)	(58.9)	(19.1)	(8.6)	(5.9)	(3.9)	(1.6)	(0.8)	(0.5)	(0.7)	(0.1)	(100.0)				
平均	79.91	79.17	77.11	76.36	76.95	74.67	71.71	78.07	75.88	64.17	79.02	78.95	79.01		
標準偏差	14.07	14.25	14.75	14.30	14.89	18.67	14.17	13.90	13.43	5.46	14.35	14.95	14.43		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査