

表 3369 透析前補正カルシウム濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	931	1,331	5,245	17,062	4,898	56,432	68,011	44,086	5,416	13,732	6,692	2,269	802	575	227,482	28,159	255,641	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.2)	(24.8)	(29.9)	(19.4)	(2.4)	(6.0)	(2.9)	(1.0)	(0.4)	(0.3)	(100.0)				
Off-line HDF	30	34	126	388	115	1,209	1,641	1,213	148	489	245	82	34	23	5,777	538	6,315	9.28	0.81
(%)	(0.5)	(0.6)	(2.2)	(6.7)	(2.0)	(20.9)	(28.4)	(21.0)	(2.6)	(8.5)	(4.2)	(1.4)	(0.6)	(0.4)	(100.0)				
On-line HDF	135	227	837	2,505	710	8,152	9,965	6,433	756	1,980	937	305	98	47	33,087	3,003	36,090	9.17	0.73
(%)	(0.4)	(0.7)	(2.5)	(7.6)	(2.1)	(24.6)	(30.1)	(19.4)	(2.3)	(6.0)	(2.8)	(0.9)	(0.3)	(0.1)	(100.0)				
Push/Pull HDF	1	4	8	29	9	128	140	93	10	30	11	1	1	1	466	71	537	9.17	0.67
(%)	(0.2)	(0.9)	(1.7)	(6.2)	(1.9)	(27.5)	(30.0)	(20.0)	(2.1)	(6.4)	(2.4)	(0.2)	(0.2)	(0.2)	(100.0)				
ハイオフィルトレーション	2	3	3	21	5	66	85	87	14	19	13	1	2	2	323	18	341	9.30	0.75
(%)	(0.6)	(0.9)	(0.9)	(6.5)	(1.5)	(20.4)	(26.3)	(26.9)	(4.3)	(5.9)	(4.0)	(0.3)	(0.6)	(0.6)	(100.0)				
血液濾過	1		1	3	1	17	19	23	2	12	8		1	1	89	3	92	9.51	0.83
(%)	(1.1)		(1.1)	(3.4)	(1.1)	(19.1)	(21.3)	(25.8)	(2.2)	(13.5)	(9.0)		(1.1)	(1.1)	(100.0)				
血液吸着透析 (リクセル)	9	11	43	113	39	334	431	313	44	129	54	11	5	4	1,540	83	1,623	9.21	0.78
(%)	(0.6)	(0.7)	(2.8)	(7.3)	(2.5)	(21.7)	(28.0)	(20.3)	(2.9)	(8.4)	(3.5)	(0.7)	(0.3)	(0.3)	(100.0)				
血液吸着透析 (その他)				1		1	1								3	4	7	8.63	0.32
(%)				(33.3)		(33.3)	(33.3)								(100.0)				
在宅血液透析	4	3	9	31	9	74	109	92	17	29	14	1	1	1	394	127	521	9.22	0.81
(%)	(1.0)	(0.8)	(2.3)	(7.9)	(2.3)	(18.8)	(27.7)	(23.4)	(4.3)	(7.4)	(3.6)	(0.3)	(0.3)	(0.3)	(100.0)				
腹膜透析 (CAPD)	15	17	66	142	42	458	877	877	114	319	167	72	16	16	3,198	1,823	5,021	9.44	0.80
(%)	(0.5)	(0.5)	(2.1)	(4.4)	(1.3)	(14.3)	(27.4)	(27.4)	(3.6)	(10.0)	(5.2)	(2.3)	(0.5)	(0.5)	(100.0)				
腹膜透析 (APD)	11	14	53	100	31	325	535	519	48	157	95	30	13	9	1,940	899	2,839	9.35	0.81
(%)	(0.6)	(0.7)	(2.7)	(5.2)	(1.6)	(16.8)	(27.6)	(26.8)	(2.5)	(8.1)	(4.9)	(1.5)	(0.7)	(0.5)	(100.0)				
腹膜透析 (CCPD)	3	5	8	44	13	123	190	233	29	81	34	9		2	774	307	1,081	9.40	0.74
(%)	(0.4)	(0.6)	(1.0)	(5.7)	(1.7)	(15.9)	(24.5)	(30.1)	(3.7)	(10.5)	(4.4)	(1.2)		(0.3)	(100.0)				
合計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし (%)																			
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査